



Supporting Online Material for
Spreading Dead Zones and Consequences for Marine Ecosystems

Robert J. Diaz* and Rutger Rosenberg

*To whom correspondence should be addressed. E-mail: diaz@vims.edu

Published 15 August 2008, *Science* **321**, 926 (2008)
DOI: 10.1126/science.1156401

This PDF file includes:

Fig. S1
Tables S1 and S2
References

Supplemental Online Material

Spreading Dead Zones and Consequence for Marine Ecosystems.

Robert J. Diaz,^{1*} Rutger Rosenberg²

¹Virginia Institute of Marine Science, College of William and Mary, Gloucester Pt., VA 23062.

²Department of Marine Ecology, University of Gothenburg, Kristineberg Marine Research Station, 450 34 Fiskebäckskil, Sweden.

*To whom correspondence should be addressed. E-mail: diaz@vims.edu

This PDF file includes

Figure S1

Tables S1 and S2

Supplemental References




Figure S1. Cumulative increase in dead zones through time reported in the scientific literature. Systems are grouped by decade of first documented account (Table S1). The number of dead zones started to approximately double every ten years starting in the 1960s.

Table S1. Hypoxic areas around the world linked to eutrophication. Hypoxia type is characterized as Episodic: events occurring at irregular intervals >1 year; Periodic: several to many events per year lasting from hours to weeks, also includes daily hypoxia; Seasonal: yearly events related mostly to summer or autumn seasons; Persistent: year-round or near year-round hypoxia. Decade of first or recent data report is an estimate of when hypoxia first started. Area is the maximum reported spatial extent of hypoxia, however, most systems do not have area estimates.

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Episodic	North Sea, SE	Denmark	1980		Mortality with annual recolonization	Stressed		Dyer et al. 1983, Westernhagen & Dethlefsen 1983
Episodic	Wismar Bay	Denmark	1980		Mortality with reduced benthos	Stressed		Prena 1995a, 1995b
Episodic	Bude Bay	England	1990	4	Mortality along 12 km of coastline	Die-off of <i>Nuccella lapillus</i> and <i>Cenebra erinacea</i>	In vicinity of sewage outfall	Gibbs et al. 1999
Episodic	Gulf of Riga	Estonia, Latvia	1980					Karlsou et al. 2002, Ojaveera & Andrushaitis 2004
Episodic	Baie de Somme	France	1980		Mass Mortality with multi-year recovery	Collapse of cockle fishery	Eutrophication first became apparent in 1982-85	Desprez et al. 1992 Rybarczyk et al. 1996
Episodic	Bay of Vilaine	France	1990					Le Bris & Glemarec 1995
Episodic	Hohwacht Bay	Germany	1910			Fish killed in nets	Combination of anthropogenic and natural factors.	Gerlach 1990
Episodic	Kalifornien & Schoenberger Strand	Germany	1920			Fish kills	Combination of anthropogenic and natural factors.	Gerlach 1990
Episodic	Donegal Bay	Ireland	2000				low DO from HAB	Silke et al. 2005
Episodic	Tokyo Bay	Japan	1970	683	Mass Mortality	Decline of Mantis shrimp	55-65% of Bay hypoxic July-August	Kodama et al. 2002, 2006
Episodic	Mauritania Coast	Mauritania	1990				Anoxia related to severe dry climate conditions	Le Loeuff 1999
Episodic	East Frisian, Wadden Sea	Netherlands	1990				Debated as to natural or not	Kaiser & Lutter 1998
Episodic	Wadden Sea	Netherlands	1990	3000	Mortality	Decrease in percentage of filter feeders in benthic biomass	Increased macroalgae	de Jonge et al. 1994
Episodic	Cape Rodney, near	New Zealand	1980	8000		Mortality	HAB related to clam weather caused low DO and anoxia	Taylor et al. 1985
Episodic	Paracas Bay	Peru	2000			Mortality	Harmful algal bloom	Kahru et al. 2004
Episodic	Manila Bay	Phillipines	1980				Sewage discharge	Velasquez et al. 2002
Episodic	Volga River Delta	Russia	1990				Harmful algal blooms	Bukharitsin & Luneva 1994
Episodic	Loch Ailort	Scotland	1990		Reduced		Salmon farms	Gillibrand et al. 1996
Episodic	Gullmarsfjord, Alsback Deep	Sweden	1980	25	Mass Mortality with multi-year recovery	Low foram diversity at deep periodically hypoxic/anoxic areas	Hypoxia evolves in autumn and winter.	Gustafsson & Nordberg 2001, Nilsson & Rosenberg 2000
Episodic	Pak Panang Bay	Thailand	2000				Combination of pollution and weather	Foulkes et al. 2007
Episodic	Alamitos Bay	US-California	2000	2				Rabalais 1998

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Episodic	Coyote Creek	US-California	1970	5		Fisherman reported absence of fish and pelagic invertebrates, fish returned after hypoxia ended	Sewage spill	Cloern & Oremland 1983
Episodic	North San Francisco Bay Estuary	US-California	2000			Fish Kill	Anoxia develops	Lehman et al. 2004
Episodic	San Joaquin River	US-California	1960				Sewage discharge	Lehman et al. 2004
Episodic	Connecticut River	US-Connecticut	1990	12			Episodic events in tidal freshwater zone	Bricker et al. 1999
Episodic	Charlotte Harbor	US-Florida	1980	90			Improvements offset by increased nutrients and hurricane disturbance	Turner et al. 2006, Tomasko et al 2006
Episodic	Looe Key	US-Florida	1980		Mortality	Coral mortality	Rainfall events followed by low DO in sea grass and coral reef areas	Lapointe & Matzie 1996 Bricker et al. 2007
Episodic	Tampa Bay near Beacon Key	US-Florida	1990	125		SAV recovering	Improved from nutrient management	Leverone 1995, Lowery 1998, Bricker et al. 2007
Episodic	Lake Pontchartrain	US-Louisiana	1990	421	Reduced	Loss of large clams		Abadie & Poirrier 2000
Episodic	Kennebec & Androscoggin Rivers	US-Maine	2000	56				Bricker et al. 1999
Episodic	Buzzards Bay	US-Massachusetts	1990	3			Summer hypoxia	Bricker et al. 1999
Episodic	Dorchester Bay	US-Massachusetts	1990				Improved from nutrient management	Maciolek et al. 2005
Episodic	Shellbank Basin	US-New York	1990					Rhoads et al. 2001
Episodic	New York Bight	US-New York/ New Jersey	1970	987	Mass Mortality with multi-year recovery	Surf clam/finfish mortality, Avoidance, northward migration of bluefish blocked	Clam weather lead to stratification and bloom	Garlo et al. 1979, Sindermann & Swanson 1980
Episodic	Cape Fear River	US-North Carolina	1990		Reduced	Fish kills		Mallin et al. 1999, Posey et al. 1999
Episodic	Futch Creek	US-North Carolina	2000					MacPherson et al. 2007
Episodic	Hewletts Creek	US-North Carolina	2000				Heavily developed watershed	MacPherson et al. 2007
Episodic	Pamlico Sound	US-North Carolina	1990			Common finfish species (pinfish, spot, croaker) had skin lesions and signs of systemic bacterial infections	Hurricane increased freshwater and nutrient input.	Paerl et al. 2000
Episodic	Long Bay	US-South Carolina	2000				High Cholera	Koepfner et al. 2007
Episodic	Baffin Bay	US-Texas	1980	61				Bricker et al. 1999
Episodic	Bryan Mound, 10 m	US-Texas	1970		Mass Mortality with multi-year recovery	Stressed		Harper et al. 1981, 1991
Episodic	Gulf of Mexico, Freeport	US-Texas	1970	50	Mortality with multi-year recovery	Avoidance, some mortality.		Harper & Rabalais 1995
Episodic	Lavaca & Chocolate Bay	US-Texas	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Episodic	Bryan Mound, 20 m	US-Texas	1980		Mortality with annual recolonization	Decreased diversity and abundance at the onset of hypoxia		Harper et al. 1981, 1991

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Episodic	Morrocoy National Park	Venezuela	1990				Climate and nutrients lead to bloom, resulted in decline in coral cover from 43% to 5%	Isaza et al. 2006
Episodic	Krka Estuary	Yugoslavia	1960		Mass Mortality	Mortality	low DO first reported in 1969	Legovic et al. 1991
Periodic	Harvey Estuary	Australia	1980				Associated with bloom	Hearn & Robson 2001
Periodic	Patos Lagoon	Brazil	2000			Fish kills	Combination of anthropogenic and natural factors.	Marques et al. 2004
Periodic	Gironde Estuary	France	1990					Abril et al. 1999
Periodic	Prevost Lagoon	France	1990		Mass Mortality with annual recolonization	Reduced aquaculture production	Increased macroalgae	Guyoneaud et al. 1998
Periodic	Chinhae Bay	Korea	1970	185		Reduced		Lim et al. 2006
Periodic	Westerschelde Estuary	Netherlands	1990				Industry and live stock farming, DO increase during day produce feedback mechanisms affects denitrification/nitrification.	Irigoiien et al. 1999
Periodic	Ria Formosa	Portugal	2000					Nobre et al. 2005
Periodic	Jobos Bay	Puerto Rico	1990					Sanger et al. 2002
Periodic	Koljo Fjord	Sweden	1940		Mortality with annual recolonization	Low foram diversity at deep periodically hypoxic/anoxic areas, low macrofaunal biomass	Periodic hypoxia has been noted sine the 1940s	Gustafsson & Nordberg 1999, Rosenberg et al. 2001
Periodic	Bon Secour Bay	US-Alabama	2000		Mortality	Loss of oyster		Rikard et al. 2000
Periodic	Newport Bay	US-California	1990	0.8				Rabalais 1998
Periodic	San Diego Bay	US-California	1990	4				Rabalais 1998
Periodic	Tijuana Estuary	US-California	2000	0.1				Bricker et al. 1999, Sanger et al. 2002
Periodic	Upper Pond, Elkhorn Slough	US-California	2000	0.4				Bricker et al. 1999
Periodic	Blackwater Landing	US-Delaware	1990					Sanger et al. 2002
Periodic	Pepper Creek	US-Delaware	2000			Fish migrate in and out of creek depending on DO	Daily cycle of DO from supersaturated to hypoxic	Tyler 2004
Periodic	Apalachee Bay	US-Florida	2000	61				Bricker et al. 1999
Periodic	Apalachicola Bay	US-Florida	1990	86				Lowery 1998, Sanger et al. 2002
Periodic	Bayou Chico	US-Florida	1990					Summers et al. 1997
Periodic	Escambia Bay	US-Florida	1990					Summers et al. 1997
Periodic	Florida Bay	US-Florida	2000	2287				Bricker et al. 1999
Periodic	Indian River	US-Florida	2000	253				Bricker et al. 1999
Periodic	Patch Reef	US-Florida	1990		Mortality		Rainfall events followed by low DO in seagrass and coral reef areas	Lapointe & Matzie 1996
Periodic	Pine Channel	US-Florida	1990		Mortality		Rainfall events followed by low DO in seagrass and coral reef areas	Lapointe & Matzie 1996
Periodic	Port Pine	US-Florida	1990		Mortality		Rainfall events followed by low DO in seagrass and coral reef areas	Lapointe & Matzie 1996

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Periodic	Sarasota Bay	US-Florida	2000	33		SAV recovering	Improved from nutrient management	Bricker et al. 1999
Periodic	St. Andrew Bay	US-Florida	2000	34				Bricker et al. 1999
Periodic	St. Joseph Bay	US-Florida	1990		Brittlestar migration	Avoidance		Leonard & McClintock 1999
Periodic	St. Lucie River	US-Florida	1990					Chamberlain & Hayward 1996
Periodic	Ten Thousand Islands North	US-Florida	2000	112				Bricker et al. 1999
Periodic	Ten Thousand Islands South	US-Florida	2000	203				Bricker et al. 1999
Periodic	Sapelo Island	US-Georgia	1990				Daily cycle of DO from supersaturated to hypoxic	Sanger et al. 2002
Periodic	Amite River	US-Louisiana	1990	3	Reduced	Loss of large clams	Part of Lake Pontchartrain.	Bricker et al. 1999
Periodic	Atchafalaya & Vermillion Bays	US-Louisiana	2000	112			Stratification highly significant factor	Bricker et al. 1999
Periodic	Breton & Chandeleur Sounds	US-Louisiana	2000	570				Bricker et al. 1999
Periodic	Calcasieu Lake	US-Louisiana	2000	129				Bricker et al. 1999
Periodic	Wells Inlet	US-Maine	1990				Daily cycle of DO from supersaturated to hypoxic, Submerged aquatic vegetation bed	Sanger et al. 2002
Periodic	Chester River	US-Maryland	1990	34	Mortality with annual recolonization	Avoidance, some mortality	Low DO events occur in June-Aug	Bricker et al. 1999
Periodic	Choptank River	US-Maryland	1990	69			Low DO events occur in June-Sept	Bricker et al. 1999
Periodic	Patuxent River	US-Maryland	1930	58	Mortality with annual recolonization	Avoidance, low egg hatching/larval mortality, bay anchovy eggs killed by hypoxia	Duration and spatial extent of low DO has increased from 1930-79	Keister et al. 2000, Breitbart et al 1997
Periodic	St. Leonard Creek	US-Maryland	1990					Summers et al. 1997
Periodic	Boston Harbor	US-Massachusetts	1980	0.8			Improved from nutrient management	Bricker et al. 1999
Periodic	Cape Cod Bay	US-Massachusetts	1990	12			Blooms	Rabalais 1998
Periodic	Waquoit Bay	US-Massachusetts	1990				Daily cycle of DO from saturated to hypoxic	Fritz et al. 1996, D'Avanzo & Kremer 1994
Periodic	East Mississippi Sound	US-Mississippi	2000	181			Stratification highly significant factor	Bricker et al. 1999
Periodic	West Mississippi Sound	US-Mississippi	2000	266				Bricker et al. 1999
Periodic	Berger Basin	US-New York	1970					Rhoads et al. 2001
Periodic	Grassy Bay	US-New York	1970	26			Areas anoxic in June-Aug	Rhoads et al. 2001, Bricker et al. 1999
Periodic	Great South Bay	US-New York	1990	30			Improved, but tidal creeks periodically hypoxic between July and September	Bricker et al. 1999
Periodic	New River	US-North Carolina	2000	21				Bricker et al. 1999

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Periodic	Pages Creek	US-North Carolina	2000					MacPherson et al. 2007
Periodic	South Slough	US-Oregon	1990				Daily cycle of DO from supersaturated to hypoxic	Sanger et al. 2002
Periodic	Narragansett Bay	US-Rhode Island	1950	93			Hypoxic June to September	Bergondo et al. 1950, Deacutis et al. 2006, Bricker et al. 1999
Periodic	Beresford Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Broad River	US-South Carolina	2000	70				Bricker et al. 1999
Periodic	Bull Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Charleston Harbor	US-South Carolina	2000	33			improved wastewater treatment, phosphate ban.	Bricker et al. 1999
Periodic	Deep Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Diesel Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Foster Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	James Island Creek	US-South Carolina	1990		Mortality/avoidance	Avoidance		Burnett 1997
Periodic	Malind Creek	US-South Carolina	2000					Gillett et al. 2005
Periodic	New Market Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Okatee Creek	US-South Carolina	2000					Gillett et al. 2005
Periodic	Orange Grove Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Rathall Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	Shem Creek	US-South Carolina	1990					Lerberg et al. 2000
Periodic	St. Helena Sound	US-South Carolina	2000	71				Bricker et al. 1999
Periodic	Winyah Bay	US-South Carolina	1990	29				Bricker et al. 1999, 2007
Periodic	Savannah River	US-South Carolina/Georgia	2000	23			Nonpoint sources a factor	Bricker et al. 1999
Periodic	Brazos River	US-Texas	2000	22				Bricker et al. 1999
Periodic	Upper Laguna Madre	US-Texas	2000	293				Bricker et al. 1999
Periodic	Goodwin Island	US-Virginia	1990				Submerged aquatic vegetation bed	Sanger et al. 2002
Periodic	Onancock Creek	US-Virginia	2000					Wang 2005
Periodic	Rappahannock River	US-Virginia	1990	92	Mortality with annual recolonization	Avoidance	Some anoxia in summer	Llanso' 1992
Periodic	Taskinas Creek	US-Virginia	1990					Sanger et al. 2002
Periodic	York River	US-Virginia	1980	43	No response	Avoidance, reduced diversity following low DO event	7 to 8 hypoxic events between June and Sept	Pihl et al. 1991, Diaz et al. 1992, Sagasti et al. 2001
Periodic	South Puget Sound	US-Washington	2000	34				Bricker et al. 1999
Periodic	Whidbey Basin & Skagit Bay	US-Washington	1990	26				Rabalais 1998
Periodic/Seasonal	Palude della Rosa	Italy	1990		Mortality with annual recolonization		Increased macroalgae	Tagliapietra et al. 1998
Periodic/Seasonal	Elkhorn Slough	US-California	1990	1.3				Bricker et al. 1999, Sanger et al. 2002

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Periodic/Seasonal	Saco Bay	US-Maine	2000	5				Bricker et al. 1999
Periodic/Seasonal	Great Bay	US-New Hampshire	2000	0.6				Bricker et al. 1999
Seasonal	Cabbage Tree Basin	Australia	1980		Mortality with annual recolonization			Rainer & Fitzhardinge 1981
Seasonal	Horseshoe Lagoon	Australia	1990					Donnelly et al. 1999
Seasonal	Chetumal Bay	Belize	2000			Coral reef affected	Untreated sewage, agricultural runoff	Isaza et al. 2006
Seasonal	Conceição Lagoon	Brazil	1980			Fish kills	Anoxia develops	Fonseca 2006
Seasonal	Guanabara Bay	Brazil	1990		Mortality		Tropical	Valentin et al. 1999
Seasonal	Pina Basin	Brazil	1990					Somerfield et al 2003
Seasonal	Rodrigo de Freitas Lagoon	Brazil	1980				Anoxia develops	Marques et al. 2004
Seasonal	Saanich Inlet	Canada	1960					Tunncliffe 1981
Seasonal	Tracadie & Winter Bays	Canada	2000				Fish farming	Miron et al. 2005
Seasonal	East China Sea	China	1950	15000			Discharge from Yangtze River is source of nutrients	Li & Daler 2004, Wang 2006, Chen et al. 2007
Seasonal	Tolo Harbor	China	1980	80	Mass mortality	Mass mortality	Rapid (several months to one year) winter recovery of fauna	Wu 1982
Seasonal	Als	Denmark	1920	40				Gerlach 1990
Seasonal	Århus Bay	Denmark	1980	1300	Mass Mortality with multi-year recovery	<i>Nephtys hombergi</i> (more low DO tolerant) displaced <i>N. ciliata</i> after hypoxic event	low DO event killed most of macrofauna; macrofauna recolonized following year	Fallesen & Jørgensen 1991, Josefson & Hansen 2004
Seasonal	Belt Sea	Denmark	1970	2150				Karlson et al. 2002
Seasonal	Coastal waters, Denmark	Denmark	2000	9000			2001 worse hypoxia year recorded, high runoff, low wind	Conley et al. 2007
Seasonal	Flensborg Fjord	Denmark	1980					Josefson & Hansen 2004
Seasonal	Kattegat (NW)	Denmark	1980					Josefson & Hansen 2004
Seasonal	Kattegat (S)	Denmark	1980					Josefson & Hansen 2004
Seasonal	Køge Bugt	Denmark	1980					Josefson & Hansen 2004
Seasonal	Lillebælt N	Denmark	1980					Josefson & Hansen 2004
Seasonal	Limfjord/Løgstør Bredn.	Denmark	1980	440	Mass Mortality with annual recolonization	Demersal fisheries gone.		Jørgensen 1980, Hylleberg 1993
Seasonal	Little Belt	Denmark	1910	450				Gerlach 1990
Seasonal	Mariager Fjord	Denmark	1990	20				Fallesen et al. 2000
Seasonal	Nissum Fjord Inner & Middle	Denmark	1980					Josefson & Hansen 2004
Seasonal	Nivå Bugt	Denmark	1980					Josefson & Hansen 2004

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference	
Seasonal	Odense Fjord Outer Basin	Denmark	1980					Josefson & Hansen 2004	
Seasonal	Ringgårdsbassin	Denmark	1980					Josefson & Hansen 2004	
Seasonal	Ringkøbing Fjord	Denmark	1980					Josefson & Hansen 2004	
Seasonal	Roskilde Fjord	Denmark	1990				85% increase in Nitrogen input from 1950s to 1990s, mainly from fertilizer	Clarke et al. 2003, Flindt et al. 1997	
Seasonal	Roskildefjord North	Denmark	1980					Josefson & Hansen 2004	
Seasonal	Skagerrak Coast	Denmark	1980				Transboundary transports of nutrients.	OSPAR 2003	
Seasonal	Skive Fjord	Denmark	1990					Josefson & Hansen 2004, Conley et al. 2007	
Seasonal	Vejle Fjord	Denmark	1980					Josefson & Hansen 2004	
Seasonal	Alexandria Harbour	Egypt	1980				Many stressors, maritime activities, agriculture, stored fertilizers, domestic waste, industrial waste	Dorgham et al. 2004	
Seasonal	Forth Estuary	England	1960			Fish kills	Improved from nutrient management	Balls et al. 1996, Griffiths 1987	
Seasonal	Mersey Estuary	England	1850	25		With recovery fish returned, Salmon runs from 1990s	Improved from nutrient management, Still has annual hypoxia in one spot upriver	Jones 2006	
Seasonal	Thames Estuary	England	1920				Worse 1930-1970, Improved from nutrient management	Andrews & Rickard 1980, Araujo et al. 1999	
Seasonal	Åland archipelago	Finland	1990	70	Mortality with reduced benthos	Reduced abundance, biomass, diversity under algal mats	Algae has increased in this area since the early 1990s	Norkko & Bonsdorff 1996, Karlson et al. 2002	
Seasonal	Archipelago Sea	Finland	1990				Fish farming	Kraufvelin et al. 2001	
Seasonal	Inre Verkviiken	Finland	1970	0.5			Water below 10-15 m becomes anoxic in late summer, but the thickness of the anoxic zone varies year to year	Lindholm 1996	
Seasonal	Neva Bay	Finland	1990	300				Karlson et al. 2002	
Seasonal	Etang de Berre	France	1970	132				Hydraulic modifications (dam)	Stora & Arnoux 1983
Seasonal	Loire Estuary	France	1990					Mortality of migratory species, mortality of mullet <i>Liza ramada</i>	Thouvenin et al. 1994, Abril et al. 2003
Seasonal	Seine Estuary	France	1990					DO deficiency increasing	Garnier et al. 2001
Seasonal	Thau Lagoon	France	1990		Mass Mortality	Mortality/Reduced shellfish production	Semi-intensive aquaculture farming	Mazouni et al. 1996, Chapelle et al. 2000, Souchu et al. 1998	
Seasonal	Eckernforde Bay	Germany	1910		Mortality with annual recolonization	Mass mortalities 1875, 1931, 1936-37	Benthic community annually goes through a boom-and-crash cycle	Gerlach 1990, Brongersma-Sanders 1957, D'Andrea et al. 1996, Bentley & Nittrouer 1999	
Seasonal	Elbe Estuary	Germany	1980			Larval and juvenile fishes less able to escape low DO and result in mortality	Improved from nutrient management	Thiel et al. 1995	

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Seasonal	German Bight	Germany	1980	15000	Mass Mortality with annual recolonization	Mortality		Dethlefsen & Westernhagen 1983, Brockmann et al. 1988, Niermann et al. 1990
Seasonal	Kiel Bay	Germany	1960	890	Mass Mortality with annual recolonization	Stressed		Arntz 1981, Rumohr 1986, Weigelt 1990 1991, Oeschger & Storey 1990
Seasonal	Lueback Bay	Germany	1960					Gerlach 1990
Seasonal	Mecklenburg Bay	Germany	1980	1860				Karlson et al. 2002
Seasonal	Pomeranian Bay	Germany	1990	390	Mass Mortality with multi-year recovery	Amphipods disappeared from hypoxic zones	low DO 1st observed in 1994	Powilleit & Kube 1999, Karlson et al. 2002
Seasonal	Oder Lagoon	Germany, Poland	1990					Pohl et al. 1998
Seasonal	Elefsis Bay	Greece	1980	67	Mass Mortality with annual recolonization		Low DO first observed in 1973	Friligos & Zenetos 1988, Theodorou 1996
Seasonal	Gialova Lagoon	Greece	1990	2500	Mortality with annual recolonization		Limited exchange with adjacent lagoon	Arvanitidis et al. 1999
Seasonal	Lough Ine	Ireland	1970		Mass Mortality with annual recolonization			Kitching et al. 1976
Seasonal	Gulf of Aqaba	Israel	1990			High coral mortality	Fish farming	Eden et al. 2003, Erez et al. 2004
Seasonal	Adriatic Sea, northern	Italy	1960	3750	1969 - 1st mass mortality of benthos		Seasonal anoxia increased in 1960; reached maximum in 1981-1987	Justic' et al 1987, 1993, Barmawidjaja et al. 1995
Seasonal	Goro Lagoon	Italy	1990		Mortality with annual recolonization			Reizopoulou et al. 1996
Seasonal	Orbetello Lagoon	Italy	1990					Lardicci et al. 2001
Seasonal	Venice Lagoon	Italy	1990					Flindt et al. 1997
Seasonal	Gulf of Trieste	Italy, Slovenia, Croatia	1960		Mass Mortality with multi-year recovery	Stressed	Mass mortality recorded in 1974	Stachowitsch 1984, 1991, Simunovic et al. 1999, Justic' et al 1987, Fedra et al. 1976
Seasonal	Ariake & Isahaya Bays	Japan	2000	240			Hydraulic modifications (dam)	Hodoki & Murakami 2006
Seasonal	Beppu Bay	Japan	1970				Enclosed topography	Suzuki 2001
Seasonal	Dokai Bay	Japan	1990	11	Mortality		Low DO is upwelled into shallower areas by winds	Ueda et al. 2000
Seasonal	Funka Bay	Japan	1990			Avoidance		Kimura et al. 2004
Seasonal	Gokasho Bay	Japan	1990	1.0			Combination of anthropogenic (fish farming) and natural factors	Yokoyama 2002
Seasonal	Hakata Bay	Japan	1970	120	Mortality with annual recolonization			Karim et al. 2002
Seasonal	Hiuchi Sound	Japan	1970		Mass Mortality	All fauna dead at DO = 0.4 ppm		Sanukida et al. 1984
Seasonal	Ise Bay	Japan	1990	1740	Mortality	Shellfish and fisheries in shallow areas and tidal flats reduced	low DO is upwelled into shallower areas by winds and causes damage.	Nakata et al. 1997
Seasonal	Kumihama Bay	Japan	1980					Yokoyama 1995

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Seasonal	Lake Nakaumi	Japan	1990			Mortality/avoidance		Kasai et al. 2005, Yamamuro et al. 2000
Seasonal	Lake Shinji	Japan	1990		Mass Mortality			Yamamuro et al. 1998
Seasonal	Mikawa Bay	Japan	1980	500	Mortality/avoidance		low DO occurs in summer	Suzuki & Matsukawa 1987, Suzuki 2001
Seasonal	Omura Bay	Japan	1980	100			Enclosed topography	Iizuka & Min 1989
Seasonal	Osaka Bay	Japan	1980		Recruitment on increasing DO		Industrial and domestic discharges	Yokoyama 1995, Suzuki 2001
Seasonal	Seto Inland Sea	Japan	1980		Mortality	Mortality		Imabayashi 1986
Seasonal	Suo-Nada	Japan	1990					Suzuki 2001
Seasonal	Tome Cove	Japan	1980					Tsutsumi 1987
Seasonal	Chonsu Bay	Korea	2000			Reduced		Lim et al. 2006
Seasonal	Kamak Bay	Korea	1900			Reduced		Lim et al. 2006
Seasonal	Jinhae Bay	Korea	2000			Oyster mortality related to DO and red tide	Related to summer algal blooms, annual from 1980s	Lee & Kim 2008
Seasonal	Masan Bay	Korea	2000				Related to summer algal blooms	Lee & Kim 2008
Seasonal	Youngsan Estuary	Korea	2000	3		Reduced		Lim et al. 2006
Seasonal	Bojorquez Lagoon	Mexico	1980				Sewage discharge from city of Cancun	Valdes-Lozano et al. 2006
Seasonal	Oslofjord	Norway	1910	150	Mortality with annual recolonization	Reduced		Petersen 1915, Mirza & Gray 1981, Rosenberg et al. 1987
Seasonal	Bilbao Estuary	Spain	1990		Mortality	Some fauna survived with reduced activity		Gonzales-Oreja & Saiz-Salinas 1998, Saiz-Salinas & Frances-Zubillaga 1997
Seasonal	La Coruna Bay	Spain	1990					Lopez-Jamar et al. 1995
Seasonal	Ria San Martin	Spain	2000				Combination of sewage, pulp mill, industry, treatment plant being built by Torrelavega	Puente et al. 2007
Seasonal	Gullmarsfjord, Kilviken	Sweden	1970		Decreased species richness, abundance			Josefson & Widbom 1988, Hendelberg & Jensen 1993
Seasonal	Havstens Fjord	Sweden	1960		Low foram diversity and abundance at deeper hypoxic/anoxic areas		Annual hypoxia or anoxia recorded since 1962	Gustafsson & Nordberg 2000
Seasonal	Laholm Bay	Sweden	1980		Mortality with annual recolonization	Mortality		Baden et al. 1990b, Rosenberg & Loo 1988
Seasonal	Skagerrak Coast	Sweden	1950		Mortality	Stressed	Linear decrease in DO since the 1960s	Johannessen & Dahl 1996a, 1996b
Seasonal	St. Anna Archipelago	Sweden	1980	25				Karlson et al. 2002
Seasonal	Swedish West Coast Fjords	Sweden	1980			Stressed		Josefsen & Rosenberg 1988
Seasonal	Kattegat (SE)	Sweden, Denmark	1980	3850	Mass Mortality with multi-year recovery	Reduced demersal fish abundance, Norway lobster collapse	DO lowest in Sept-Oct	Petersen & Pihl 1995, Baden et al. 1990a, Josefson & Jensen 1992, Rosenberg et al. 1992

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Seasonal	Bornholm Basin	Sweden, Finland	1950	6793	Mass Mortality with multi-year recovery			Tulkki 1965, Leppakoski 1969, Karlson et al. 2002
Seasonal	Prawn culture ponds	Taiwan	2000		Mortality	Mortality of prawns, reduced growth, depressed immune system in <i>Macrobrachium rosenbergii</i>	Decomposition of unconsumed food and feces	Cheng et al. 2002
Seasonal	Lake Tunis	Tunisia	1980			Reduced worm reefs, fish kills, and H ₂ S odor permeates the city of Tunis during the summer	May go back 100s of years	Kelly & Naguib 1984
Seasonal	Black Sea NW Shelf	Ukraine, Romania, Bulgaria	1960	40000	Mass Mortality with annual recolonization	Reduced, now recovering	First reported in the 1960s, Improved from nutrient management in the 1990s	Zaitsev 1993, Bakan & Buyukgungor 2000, Friedrich et al. 2002, Mee 2006
Seasonal	Montevideo Bay	Uruguay	2000					Danulat et al. 2002
Seasonal	Mobile Bay	US-Alabama	1940	1060		Occasional shoreward migration of hypoxia stressed fish and crustaceans (Jubilees)	Stratification highly significant factor	Loesch 1960, Pennock et al. 1994, May, 1973, Engle & Summers 1999
Seasonal	Los Angeles Harbor	US-California	1950		Mass Mortality with multi-year recovery		Improved from nutrient management	Reish 1955, 2000
Seasonal	Monterey Bay	US-California	2000				Combination of anthropogenic and natural factors.	Okey 2003
Seasonal	South San Francisco Bay	US-California	1960				Hypoxia eliminated with sewage treatment plants	Nichols et al. 1986 Bricker et al. 2007
Seasonal	Bald Eagle Creek	US-Delaware	2000			Fish kills	Eutrophication and limited tidal exchange with adjacent body of water	Luther et al. 2004
Seasonal	Delaware River, upper	US-Delaware	1910				low DO first recorded in 1915, Improved from nutrient management	Patrick 1988
Seasonal	Choctawhatchee Bay	US-Florida	1990	156				Lowery 1998
Seasonal	Hillsborough Bay	US-Florida	1980				Improved from nutrient management	Santos & Simon 1980, Johansson & Lewis 1992
Seasonal	Pensacola Bay	US-Florida	1990	76				Lowery 1998
Seasonal	Perdido Bay	US-Florida	1990	64			Stratification highly significant factor	Flemer et al. 1999
Seasonal	St. Johns River	US-Florida	1990		Mortality	Mortality		Mason 1998
Seasonal	Gulf of Mexico LA-TX shelf	US-Louisiana	1970	22000		Avoidance	Area coincides with historic white and brown shrimp fishing grounds	Turner & Rabalais 1994, Justic' et al. 1996, Sen Gupta et al. 1996, Rabalais et al. 2007
Seasonal	West Hackberry, Gulf of Mexico	US-Louisiana	1980		Mortality with annual recolonization			Gaston 1985
Seasonal	Chesapeake Bay Mainstem	US-Maryland	1930	2751	Mortality with annual recolonization	Killed crabs in crab pots	Seasonal anoxia detected in sediment record as far back as 1934-1948	Zimmerman & Canual 2000, Newcombe & Horne 1938, Officer et al. 1984, Seliger et al. 1985, Holland et al., 1987, Boesch et al. 2001, Hagy et al. 2004

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Seasonal	Patapsco River	US-Maryland	2000					Riedel et al. 1999
Seasonal	Potomac River	US-Maryland	1920	264	Mortality with annual recolonization		low DO in May-Sep, increased spatial coverage of hypoxia	Sale & Skinner 1917 Bricker et al. 1999
Seasonal	Dead-end canals	US-Maryland/Delaware	1990	5			Dead-end canal have limited tidal exchange with adjacent body of water, as many as 20 canals	Maxted et al. 1997
Seasonal	Charles River	US-Massachusetts	1980					Taylor 2000
Seasonal	Fore River	US-Massachusetts	1980					Neponset River Watershed Association 2004
Seasonal	Herring River	US-Massachusetts	1980			Fish kills, decline of alewife fishery	Anoxia 1-3 weeks	Portnoy 1991
Seasonal	Barnegat Inlet	US-New Jersey	1990		Mass Mortality with annual recolonization			Moser 1998
Seasonal	Great Egg Harbor River	US-New Jersey	1990					Glenn et al. 1996
Seasonal	Mullica River Estuary	US-New Jersey	1990					Glenn et al. 1996
Seasonal	Townsend-Hereford Inlet	US-New Jersey	1990					Glenn et al. 1996
Seasonal	East River	US-New York	1920				Improved from nutrient management	Parker & O'Reilly 1991
Seasonal	Flushing Bay	US-New York	1990		Mortality			Rhoads et al. 2001
Seasonal	Hudson River	US-New York	1960	65			Improved from nutrient management, no longer hypoxic	Brosnan & O'Shea 1996
Seasonal	Long Island Sound	US-New York	1970	1036		Avoidance, some mortality.	Hypoxic June to September; duration, frequency, spatial coverage all increased since 1987	Howell & Simpson 1994, Welsh et al. 1994, Schimmel et al. 1999, Anderson & Taylor 2001
Seasonal	Mill Basin	US-New York	1990			Avoidance		Rhoads et al. 2001
Seasonal	New York City Harbor	US-New York	1990		Mass Mortality with annual recolonization	Stressed	Improved from nutrient management	Parker & O'Reilly 1991
Seasonal	Norton Basin	US-New York	1990			Avoidance		Rhoads et al. 2001
Seasonal	Raritan Bay	US-New York/New Jersey	1970	55			Improved from nutrient management	Christensen & Packard 1976
Seasonal	Albemarle & Pamlico Sounds	US-North Carolina	1990	464	Mortality	Mortality	Hurricane increased freshwater and nutrient input.	Bricker et al. 1999
Seasonal	Neuse River Estuary	US-North Carolina	1990	230	Mortality/avoidance	Fish kills, mortality of oyster.		Paerl et al. 1995, 1998, Lenihan & Peterson 1998, Lenihan 1999
Seasonal	Pamlico & Pungo Rivers	US-North Carolina	1960	44	Mass Mortality, low macrobenthic diversity and density in summer, recolonization by winter	Mortality		Tenore 1972, Hobbie et al. 1975, Stanley & Nixon 1992
Seasonal	Stono & North Edisto Rivers	US-South Carolina	2000	22				Bricker et al. 1999
Seasonal	Aransas Bay	US-Texas	1990					Lowery 1998

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Seasonal	Corpus Christi Bay	US-Texas	1980	123	Reduced	Biomass at hypoxic stations is 96% less than at normoxic stations.	Low bottom DO when bottom salinity was high	Ritter & Montagna 1999, Montagna & Ritter 2006, Montagna & Kalke 1992
Seasonal	Galveston Bay	US-Texas	1990	250				Lowery 1998, Bricker et al. 1999
Seasonal	Matagorda Bay	US-Texas	1990	281				Lowery 1998
Seasonal	San Antonio Bay	US-Texas	1990	109			Stratification highly significant factor	Lowery 1998
Seasonal	Chesapeake Bay, lower	US-Virginia	1980				Lower diversity, lower biomass, lower proportion of deep-dwelling biomass at hypoxia-affected areas	Dauer et al 1992
Seasonal	Hood Canal	US-Washington	1980	238				Paulson et al. 1993, Bricker et al. 2007
Seasonal/Episodic	Delaware River	US-Delaware	1990			Recovering American shad and striped bass fishery	Improved from nutrient management	Weisberg et al. 1996, Summers et al. 1997
Seasonal/Episodic	Biscayne Bay	US-Florida	2000	2			Related to releases of freshwater from canals	Bricker et al. 1999
Seasonal/Episodic	Barataria Bay	US-Louisiana	2000	207				Bricker et al. 1999
Seasonal/Persistent	Sea of Azov	Russia, Ukraine	1930	8560	Mortality with reduced benthos	Lower production	Prone to stratification, large anoxia 1937, 1946, 1987, 2001	Balkas et al. 1990, Chechum 1998, Debolskaya et al. 2008
Seasonal/Persistent	Skagerrak coast (Fjords)	Sweden	1920	54			Increased sedimentation of phytoplankton because of decreased grazing by zooplankton	Rosenberg 1990
Seasonal?	Imboassica Lagoon	Brazil	1990				Sewage discharge	Kozlowsky-Suzuki & Bozelli 2002
Seasonal?	Alykes Kitrous Lagoon	Greece	1990		Meiofauna reduced		Eutrophic hypersaline lagoon	Doulgeraki et al. 2006
Persistent	Scheldt Estuary	Belgium	1990					Verlaan et al. 1998
Persistent	St. Lawrence Estuary	Canada	1980	1300		Avoidance	70 year record of declining DO, anoxia formed in last several decades	Gilbert et al. 2005, Benoit et al. 2006
Persistent	Pearl River Estuary	China	2000	20				Dai et al. 2006
Persistent	Gulf of Finland, Deep	Finland	1960	5700	Reduced		Has periods of improved conditions, recolonized after renewal of bottom water	Laine et al. 1997, Andersin & Sandler 1991, Karlson et al. 2002
Persistent	Big Glory Bay	New Zealand	2000		Mass Mortality		Salmon farming	Morrissey 2000
Persistent	Gdansk Basin	Poland	1960	1200				Karlson et al. 2002
Persistent	Caspian Sea	Russia, Iran, Azerbaijan, Turkmenistan	1990		Mortality with reduced benthos		Depends on discharge intensity of Volga; high-flow years cause low DO in deep areas.	Dumont 1998
Persistent	Loch Creran	Scotland	1970		Mass Mortality			Gage 1972
Persistent	Sullom Voe	Scotland	1980		Mass Mortality			Pearson & Eleftheriou 1981
Persistent	Arkona Basin	Sweden	1980	1000				Karlson et al. 2002
Persistent	Byfjord	Sweden	1970		Mortality with reduced benthos	Only pelagic species		Rosenberg 1990, Rosenberg et al 1977

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Persistent	Gotland Basin, East	Sweden	1960	39600	Mortality with reduced benthos	Low DO prevents development of cod eggs, cod prey reduced (<i>Saduria entomon</i> and <i>Lumpenus lampetraeformis</i>)	Recolonized after renewal of bottom water	Laine et al. 1997, de Jonge et al. 1994, Karlson et al. 2002
Persistent	Gotland Basin, North	Sweden	1960	14300	Mortality with reduced benthos	Avoidance, mortality/low hatch cod eggs	Recolonized after renewal of bottom water	Laine et al. 1997
Persistent	Gotland Basin, West	Sweden	1960	16100	Mortality with reduced benthos	Avoidance, mortality/low hatch cod eggs	Recolonized after renewal of bottom water	Laine et al. 1997
Persistent	Himmerfjärden	Sweden	1970	11				Savage et al. 2002
Persistent	Stockholm Inner Archipelago	Sweden	1970	60	No Benthos and no recruitment			Rosenberg & Diaz 1993
Persistent	Baltic Sea, Northern	Sweden, Finland	1970			Stressed		Andersin et al. 1978
Persistent	Idefjord	Sweden, Norway	1960	80	Mortality with reduced benthos		Industry (sulfite pulp mill), after closure fjord recovered	Rosenberg 1980
Persistent	Tan Shui Estuary	Taiwan	1990					Jeng & Han 1996
Persistent	Dubai Creek	United Arab Emirates	2000		Low diversity and abundance, Capitellids dominant		High levels of organic loading from sewage	Saunders et al. 2007
Persistent	Caloosahatchee River	US-Florida	2000	4				Bricker et al. 1999
Persistent	Lower Laguna Madre	US-Texas	2000	28			Sewage and shrimp farming	Bricker et al. 1999
Upwelling	Western Indian Shelf	India	1990				Potential eutrophication led to lower DO in OMZ and expansion of area	Naqvi et al. 2000
Upwelling	California Current System	US-Oregon	2000		Mass Mortality	Mortality	Shifting wind patterns lead to extensive shallow shelf hypoxia	Grantham et al. 2004
Unknown	La Plata River Estuary	Argentina	2000			Change in trophic structure		Nagy et al. 2002
Unknown	Belgian Shelf north of Oostend	Belgium	2000					OSPAR 2003
Unknown	Ebrie Lagoon	Cote d'Ivoire	1980		Loss of biodiversity	Fish kills, loss of trophic structure	Domestic waste and industrial waste from coastal population	Ukwe et al. 2006
Unknown	Chichester Harbour	England	2000					OSPAR 2003
Unknown	Firth of Clyde Estuary	England	1980					Bock et al. 1999
Unknown	Holes Bay	England	2000					OSPAR 2003
Unknown	Humber Estuary	England	1980					Uncles et al. 1998
Unknown	Langstone Harbour	England	2000					OSPAR 2003
Unknown	Lindisfarne NNR Area	England	2000					OSPAR 2003
Unknown	North Sea	England	2000					Ducrottoy et al 2000
Unknown	Pagham Harbour	England	2000					OSPAR 2003
Unknown	Taw Estuary	England	2000					OSPAR 2003
Unknown	Tawe Estuary	England	2000					OSPAR 2003
Unknown	Tees Estuary	England	2000					OSPAR 2003
Unknown	Tresillian & Fal Estuaries	England	2000					OSPAR 2003

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Unknown	Ythan Estuary	England	2000					OSPAR 2003
Unknown	Baie d'Arguenon	France	2000					OSPAR 2003
Unknown	Baie de Douarnenez	France	2000					OSPAR 2003
Unknown	Baie de la Frenaye	France	2000					OSPAR 2003
Unknown	Baie de Lannion	France	2000					OSPAR 2003
Unknown	Baie de Morlaix	France	2000					OSPAR 2003
Unknown	Baie de Sainte-Brieuc	France	2000					OSPAR 2003
Unknown	Bassin d'Arcachon	France	2000					OSPAR 2003
Unknown	Boulogne	France	2000					OSPAR 2003
Unknown	Calais	France	2000					OSPAR 2003
Unknown	Canche Estuary	France	2000					OSPAR 2003
Unknown	Concarneau	France	2000					OSPAR 2003
Unknown	Dunkerque	France	2000					OSPAR 2003
Unknown	Ems Estuary	Germany, Netherlands	2000					OSPAR 2003
Unknown	Weser Estuary	Germany, Netherlands	2000					OSPAR 2003
Unknown	Chemu Lagoon	Ghana	1980		Mass Mortality		Domestic waste and industrial waste from coastal population	Ukwe et al. 2006
Unknown	Fosu Lagoon	Ghana	1990					Blay & Dongdem 1996
Unknown	Korle lagoon	Ghana	1980				Domestic waste and industrial waste from coastal population	Ukwe et al. 2006
Unknown	Piges	Greece	1990		Meiofauna reduced			Doulgeraki et al. 2006
Unknown	Porto Lagos Lagoon	Greece	1980					Koutrakis et al. 2004
Unknown	Estero la Jagua	Honduras	1990			Fish kills, decline in artisan fishery of all species, lower CPUE	Shrimp farming	Ward 2000
Unknown	Iranian Bank	Iran	1980					Dumont 1998
Unknown	Bandon Estuary Lower	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Bandon Estuary Upper	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Barrow Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Blackwater Estuary Lower	Ireland	2000					OSPAR 2003
Unknown	Broadmeadow Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Cashen Feale Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Castletown Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Colligan River	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Dungarvan Harbor	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Fergus Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Killybegs Harbour	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Unknown	Lee (Tralee) Estuary Lower	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Lee (Tralee) Estuary Upper	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Lough Mahon & Lee Estuary	Ireland	1990					Toner et al. 2005
Unknown	Owennacurra Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Slaney Estuary Lower	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Upper Feale Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Upper Suir Estuary	Ireland	1990				DO deficiency increased 1995-99	OSPAR 2003
Unknown	Hiroshima Bay	Japan	1990					Kim et al. 1997
Unknown	Mauritius Island	Mauritius	1990			Coral reef affected	Agriculture (sugar cane), untreated sewage (3 dischargers)	Thomassin et al. 1998
Unknown	Dutch Western Scheldt	Netherlands	2000					OSPAR 2003
Unknown	Wester-Ems	Netherlands	2000					OSPAR 2003
Unknown	Inner Belfast Lough & Tidal Lagan	Northern Ireland	2000					OSPAR 2003
Unknown	Strangford Lough Catchment	Northern Ireland	2000					OSPAR 2003
Unknown	Arendal fjord	Norway	2000				Transboundary transport of nutrients and organic matter	OSPAR 2003
Unknown	Drammensfjord	Norway	2000				Small fjord, long residence time in deep water	OSPAR 2003
Unknown	Frierfjord	Norway	2000				Transboundary transport of nutrients and organic matter	OSPAR 2003
Unknown	Kragerøfjord	Norway	2000				Long residence time	OSPAR 2003
Unknown	Sandbukta	Norway	2000				Transboundary transport of nutrients and organic matter	OSPAR 2003
Unknown	Sandefjordsfjord	Norway	2000				Inner part hypoxic	OSPAR 2003
Unknown	Sandnesfjord	Norway	2000				Long residence time in deep water behind sills.	OSPAR 2003
Unknown	Singlefjord	Norway	2000					OSPAR 2003
Unknown	Søndeledfjord	Norway	2000				Long residence time in deep water behind sills.	OSPAR 2003
Unknown	Steindalsfjord	Norway	2000					OSPAR 2003
Unknown	Stølefjord	Norway	2000				Transboundary transport of nutrients and organic matter	OSPAR 2003
Unknown	Tønsbergfjord	Norway	2000					OSPAR 2003
Unknown	Trysfjord & Oføtfjord	Norway	1980			Fish kills during deep water renewals	Long residence time for deep water behind sills.	Dommasnes et al. 1994
Unknown	Mondego River	Portugal	1990					Flindt et al. 1997
Unknown	Ria de Aveiro Lagoon	Portugal	1980					Lopes et al 2005

Hypoxia type	System	Country/State	Decade First/Recent Data	Area (km ²)	Benthic Response	Fisheries/Other Response	Comment	Reference
Unknown	Danshuei River Estuary	Taiwan	1990				Heavy sewage load from Tipia	Lin et al. 2007
Unknown	Izmit Bay	Turkey	1990		Mass Mortality	Mass mortality	Oil spill and fire, sewage discharge	Balkis 2003
Unknown	Weeks Bay	US-Alabama	1990				Mangrove	Sanger et al. 2002
Unknown	Wolf Bay	US-Alabama	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Big Lagoon	US-Florida	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Old River	US-Florida	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Rookery Bay	US-Florida	1990				Mangrove	Sanger et al. 2002
Unknown	St. George Sound	US-Florida	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Watsons Bayou	US-Florida	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Graden Island Bay	US-Louisiana	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Grand Bay	US-Louisiana	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Wax Lake	US-Louisiana	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Jug Bay	US-Maryland	1990				Part of regional water quality monitoring program.	Sanger et al. 2002
Unknown	Back Bay of Biloxi	US-Mississippi	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Tivolie South	US-New York	1990					Sanger et al. 2002
Unknown	Masonboro Inlet	US-North Carolina	1990					Sanger et al. 2002
Unknown	Pettaquamscutt River	US-Rhode Island	1990					Wilkin & Barnes 1997
Unknown	Potters Cove	US-Rhode Island	1990					Sanger et al. 2002
Unknown	Ace Basin	US-South Carolina	1990					Sanger et al. 2002
Unknown	North Inlet-Winyah Bay	US-South Carolina	1990					Sanger et al. 2002
Unknown	Arroyo Colorado	US-Texas	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Upper Galveston Bay Channel	US-Texas	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Offatts Bayou	US-Texas	1930		Mass Mortality	Mortality	First occurrence 1936	Gunter 1942, Engle et al. 1999
Unknown	San Bernard River	US-Texas	1990				Part of regional water quality monitoring program.	Engle et al. 1999
Unknown	Padilla Bay	US-Washington	1990				Submerged aquatic vegetation bed	Sanger et al. 2002

Table S2. Approximate latitude and longitude of the hypoxic systems in Table S1. Obtained from referenced literature or Goggle Earth.

<u>System</u>	<u>Lat.</u>	<u>Long.</u>	<u>System</u>	<u>Lat.</u>	<u>Long.</u>
Ace Basin	32.56	-80.45	Bornholm Basin	55.45	19.13
Adriatic Sea, northern	44.76	12.75	Boston Harbor	42.34	-70.99
Alamitos Bay	33.75	-118.11	Boulogne	47.07	-2.09
Åland archipelago	60.28	19.82	Brazos River	28.97	-95.38
Albemarle & Pamlico Sounds	35.3	-76.11	Breton & Chandeleur Sounds	29.78	-89.07
Alexandria Harbour	31.17	29.86	Broad River	32.4	-80.72
Als	55.08	9.62	Broadmeadow Estuary	53.47	-6.17
Alykes Kitrous Lagoon	40.37	22.64	Bryan Mound, 10 m	28.9	-95.0
Amite River	30.3	-90.56	Bryan Mound, 20 m	28.37	-94.68
Apalachee Bay	29.76	-84.77	Bude Bay	50.83	-4.63
Apalachicola Bay	29.76	-84.77	Bull Creek	32.83	-80.02
Aransas Bay	28.05	-96.99	Buzzards Bay	41.7	-70.7
Archipelago Sea	60.3	22.0	Byfjord	59.03	11.22
Arendal fjord	58.46	8.77	Cabbage Tree Basin	-34.09	151.13
Århus Bay	56.04	10.56	Calais	50.96	1.85
Ariake & Isahay Bays	32.95	130.25	Calcasieu Lake	30.02	-93.31
Arkona Basin	54.94	13.83	California Current System	44.4	-124.2
Arroyo Colorado	26.36	-97.37	Caloosahatchee River	26.62	-81.9
Atchafalaya & Vermillion Bays	29.67	-91.88	Canche Estuary	50.55	1.55
Back Bay of Biloxi	30.42	-88.86	Cape Cod Bay	41.89	-70.35
Baffin Bay	27.32	-97.53	Cape Fear River	35	-76.94
Baie d'Arguenon	48.6	-2.22	Cape Rodney, near	-36.3	174.85
Baie de Douarnenez	48.17	-4.42	Cashen Feale Estuary	52.48	-9.66
Baie de la Frenaye	48.65	-2.3	Caspian Sea	41.83	51.08
Baie de Lannion	48.72	-3.57	Castletown Estuary	53.99	-6.22
Baie de Morlaix	48.73	-3.87	Charles River	42.38	-71.04
Baie de Sainte-Brieuc	48.63	-2.67	Charleston Harbor	32.91	-79.95
Baie de Somme	50.3	1.5	Charlotte Harbor	26.76	-82.13
Bald Eagle Creek	38.7	-75.1	Chemu lagoon	5.63	0.02
Baltic Sea, Northern	58.9	20.5	Chesapeake Bay Mainstem	38.74	-76.45
Bandon Estuary Lower	51.67	-8.5	Chesapeake Bay, Lower	37.82	-76.11
Bandon Estuary Upper	51.67	-8.5	Chester River	39.2	-76.1
Barataria Bay	29.52	-90.17	Chetumal Bay	18.33	-88.08
Barnegat Inlet	39.77	-74.09	Chichester Harbour	50.8	-0.92
Barrow Estuary	52.24	-6.96	Chinhae Bay	35	128.58
Bassin d'Arcachon	44.7	-1.15	Choctawhatchee Bay	30.45	-86.32
Bay of Vilaine	47.5	-2.45	Chonsu Bay	37.86	125.72
Bayou Chico	30.4	-87.26	Choptank River	38.69	-76.27
Belgian Shelf north of Oostend	51.33	3.0	Colligan River	52.1	-7.64
Belt Sea	55.5	11.0	Concarneau	47.87	-3.92
Beppu Bay	33.3	131.54	Conceição Lagoon	-27.55	-48.45
Beresford Creek	32.9	-79.89	Connecticut River	41.35	-72.3
Berger Basin	40.65	-73.82	Corpus Christi Bay	27.75	-97.29
Big Glory Bay	-46.92	168	Coyote Creek	37.47	-122.0
Big Lagoon	30.32	-87.36	Danshuei River Estuary	25.17	121.17
Bilbao Estuary	43.3	-3.0	Dead-end canals	38.52	-75.06
Biscayne Bay	25.57	-80.22	Deep Creek	32.92	-79.77
Black Sea NW Shelf	45.5	30.3	Delaware River	40.0	-75.0
Blackwater Estuary Lower	52.43	-6.32	Delaware River, upper	39.87	-75.09
Blackwater Landing	39.08	-75.46	Diesel Creek	32.82	-79.96
Bojorquez Lagoon	21.13	-86.76	Dokai Bay	33.88	130.8
Bon Secour Bay	30.3	-87.88	Donegal Bay	54.55	-8.51

<u>System</u>	<u>Lat.</u>	<u>Long.</u>	<u>System</u>	<u>Lat.</u>	<u>Long.</u>
Dorchester Bay	42.31	-71.04	Gulf of Riga	57.63	23.58
Drammensfjord	59.63	10.42	Gulf of Trieste	45.0	13.36
Dubai Creek	25.19	55.33	Gullmarsfjord, Alsback Deep	58.33	11.33
Dungarvan Harbor	52.07	-7.58	Gullmarsfjord, Kilviken	58.33	11.33
Dunkerque	51.05	2.37	Hakata Bay	33.6	130.3
Dutch Western Scheldt	51.37	3.83	Harvey Estuary	-32.67	115.67
East China Sea	31.0	124.0	Havstens Fjord	58.3	11.77
East Frisian, Wadden Sea	53.5	5.5	Herring River	41.93	-70.06
East Mississippi Sound	30.35	-88.36	Hewletts Creek	34.19	-77.87
East River	40.8	-73.92	Hillsborough Bay	27.83	-82.43
Ebrie lagoon	5.22	-4.7	Himmerfjarden	59.0	17.72
Eckernforde Bay	54.5	10.0	Hiroshima Bay	34.08	132.33
Elbe Estuary	53.8	9.1	Hiuchi Sound	34.0	133.5
Elefsis Bay	37.86	23.55	Hohwacht Bay	54.33	10.7
Elkhorn Slough	36.81	-121.78	Holes Bay	50.73	-2
Ems Estuary	53.5	7.0	Hood Canal	47.56	-123
Escambia Bay	30.55	-87.16	Horseshoe Lagoon	-36.08	146.93
Estero la Jagua	13.2	-87.4	Hudson River	41.53	-73.99
Etang de Berre	43.45	5.13	Humber Estuary	53.55	0.0
Fergus Estuary	52.7	-9.0	Idefjord	59.3	11.1
Firth of Clyde Estuary	55.7	-4.95	Imboassica Lagoon	-22.4	-41.7
Flensburg Fjord	54.83	9.79	Indian River	28.11	-80.6
Florida Bay	25.04	-80.73	Inner Belfast Lough & Tidal Lagan Impoundment	54.67	-5.6
Flushing Bay	40.63	-73.81	Inre Verkviken	60.25	20
Fore River	42.24	-70.96	Iranian Bank	37.36	53.91
Forth Estuary	56.1	-3.5	Ise Bay	34.66	136.75
Foster Creek	32.86	-79.86	Izmit Bay	40.74	28.39
Fosu Lagoon	5.12	-1.27	James Island Creek	32.76	-79.94
Frierfjord	59.1	9.62	Jinhae Bay	35.05	128.52
Funka Bay	42.33	140.67	Jobos Bay	17.96	-66.24
Futch Creek	34.3	-77.76	Jug Bay	38.45	-76.27
Galveston Bay	29.41	-94.88	Kalifornien & Schoenberger Strand	54.44	10.4
Gdansk Basin	55	19.25	Kamak Bay	34.68	127.67
German Bight	54.7	7.0	Kattegat (NW)	57.51	10.84
Gialova Lagoon	21.67	36.97	Kattegat (S)	56.14	11.52
Gironde Estuary	44.75	-1.0	Kattegat (SE)	56.86	11.46
Gokasho Bay	34.32	136.67	Kennebec & Androscoggin Rivers	44.03	-69.8
Goodwin Island	37.22	-76.4	Kiel Bay	54.37	10.53
Goro Lagoon	44.8	12.4	Killybegs Harbour	54.63	-8.43
Gotland Basin, East	57.5	20.1	Køge Bugt	55.5	12.36
Gotland Basin, North	57.95	19.96	Koljo Fjord	58.25	11.58
Gotland Basin, West	58.0	18.0	Korle lagoon	5.55	-0.22
Graden Island Bay	29.04	-89.12	Kragerøfjord	58.85	9.47
Grand Bay	29.37	-89.37	Krka Estuary	15.83	43.5
Grassy Bay	40.63	-73.79	Kumihama Bay	35.63	134.91
Great Bay	43.07	-70.87	La Coruña Bay	43.4	-8.4
Great Egg Harbor River	39.3	-74.63	La Plata River Estuary	-35.0	-57.0
Great South Bay	40.71	-73.05	Laholm Bay	56.54	12.81
Guanabara Bay	-23.0	-43.0	Lake Nakauri	35.47	133.2
Gulf of Aqaba	29.54	34.97	Lake Pontchartrain	30.1	-90
Gulf of Finland, Deep	59.82	25.5	Lake Shinji	35.5	133
Gulf of Mexico LA-TX shelf	28.83	-91.41	Lake Tunis	36.82	10.25
Gulf of Mexico, Freeport	28.95	-95.31			

<u>System</u>	<u>Lat.</u>	<u>Long.</u>	<u>System</u>	<u>Lat.</u>	<u>Long.</u>
Langstone Harbour	50.82	-1	Oder Lagoon	54.5	14.3
Lavaca & Chocolate Bay	28.58	-98.61	Offatts Bayou	29.28	-94.85
Lee (Tralee) Estuary Lower	52.28	-9.93	Okatee Creek	32.31	-80.93
Lee (Tralee) Estuary Upper	52.28	-9.93	Old River	29.8	-85.03
Lillebælt N	55.66	10.1	Omura Bay	32.97	129.88
Limfjord/Løgstør Bredn.	56.94	9.07	Onancock Creek	37.71	-75.76
Lindisfarne NNR Area	55.68	-1.78	Orange Grove Creek	32.81	-79.98
Little Belt	55.0	10.08	Orbetello Lagoon	42.44	11.2
Loch Ailort	56.25	-5.74	Osaka Bay	34.65	136.45
Loch Creran	57.3	-5.5	Oslofjord	59.4	10.72
Loire Estuary	47.2	-1.9	Owennacurra Estuary	51.87	-8.2
Long Bay	33.62	-78.88	Padilla Bay	48.52	-122.53
Long Island Sound	41.08	-72.88	Pages Creek	34.28	-77.79
Looe Key	24.56	-81.41	Pagham Harbour	50.77	-0.75
Los Angeles Harbor	33.74	-118.23	Pak Panang Bay	8.27	100.07
Lough Ine	55.56	-4.9	Palude della Rosa	45.5	12.42
Lough Mahon & Lee Estuary	51.88	-8.35	Pamlico & Pungo Rivers	35.42	-76.77
Lower Laguna Madre	26.42	-97.36	Pamlico Sound	35.4	-75.7
Lueback Bay	54.03	10.88	Paracas Bay	-14.0	-76.2
Malind Creek	32.32	-80.92	Patapsco River	39.2	-76.5
Manila Bay	14.52	120.77	Patch Reef	24.62	-81.4
Mariager Fjord	56.65	9.95	Patos Lagoon	-31.1	-51.25
Masan Bay	35.17	128.59	Patuxent River	38.38	-76.51
Masonboro Inlet	34.16	-77.85	Pearl River Estuary	23.0	113.6
Matagorda Bay	28.5	-96.4	Pensacola Bay	30.46	-86.97
Mauritania Coast	20.0	-17.0	Pepper Creek	38.56	-75.2
Mauritius Island	-20.25	57.5	Perdido Bay	30.38	-87.44
Mecklenburg Bay	54.2	11.2	Pettaquamscutt River	41.46	-71.45
Mersey Estuary	53.35	-2.9	Piges	40.87	24.65
Mikawa Bay	34.61	137.05	Pina Basin	-8.08	-34.9
Mill Basin	40.6	-73.91	Pine Channel	24.7	-81.41
Mobile Bay	30.46	-88.0	Pomeranian Bay	54.3	14.0
Mondego River	40.17	-8.65	Port Pine	24.73	-81.41
Monterey Bay	36.79	-121.9	Porto Lagos Lagoon	41.01	25.14
Montevideo Bay	-34.92	-56.22	Potomac River	38.4	-77.09
Morrocroy National Park	10.82	-68.27	Potters Cove	41.58	-71.32
Mullica River Estuary	39.5	-74.33	Prawn culture ponds	22.0	121.0
Narragansett Bay	41.68	-71.33	Prevost Lagoon	43.5	3.9
Neuse River Estuary	35.04	-76.58	Rappahannock River	37.6	-76.35
Neva Bay	60.09	29.57	Raritan Bay	40.48	-74.2
New Market Creek	32.82	-79.94	Rathall Creek	32.86	-79.89
New River	34.64	-77.34	Ria de Aveiro Lagoon	40.68	-8.65
New York Bight	39.96	-73.56	Ria Formosa	37.03	-7.78
New York City Harbor	40.67	-73.98	Ria San Martin	43.4	-4.04
Newport Bay	33.62	-117.89	Ringgårdsbassin	55.03	10.44
Nissum Fjord Inner & Middle	56.31	8.26	Ringkøbing Fjord	56.05	8.21
Nivå Bugt	56.02	12.62	Rodrigo de Freitas Lagoon	-23.98	-46.32
North Inlet-Winyah Bay	33.35	-79.19	Rookery Bay	26.05	-81.73
North San Francisco Bay Estuary	37.63	-122.18	Roskilde Fjord	55.8	12.0
North Sea	58.25	11.45	Roskildefjord North	55.95	11.98
North Sea, SE	56.1	7.2	Saanich Inlet	49.07	-123.53
Norton Basin	40.6	-73.77	Saco Bay	43.5	-70.38
Odense Fjord Outer Basin	55.51	10.55	San Antonio Bay	28.3	-96.7

<u>System</u>	<u>Lat.</u>	<u>Long.</u>	<u>System</u>	<u>Lat.</u>	<u>Long.</u>
San Bernard River	28.88	-95.47	Tijuana Estuary	32.56	-117.13
San Diego Bay	32.67	-117.14	Tivolie South	42.03	-73.92
San Joaquin River	38.05	-121.7	Tokyo Bay	35.93	139.92
Sandebukta	59.54	10.27	Tolo Harbor	20.07	110.3
Sandefjordsfjord	59.03	10.27	Tome Cove	20	110.0
Sandnesfjord	58.7	9.22	Tønsbergfjord	59.2	10.35
Sapelo Island	31.41	-81.3	Townsend-Hereford Inlet	39.12	-74.71
Sarasota Bay	27.32	-82.57	Tracadie & Winter Bays	46.39	-62.99
Savannah River	32.14	-80.94	Tresillian & Fal Estuaries	50.17	-5.03
Scheldt Estuary	51.25	4.17	Trysfjord & Ofotfjord	58.05	7.7
Sea of Azov	46.2	37.0	Upper Feale Estuary	52.44	-9.62
Seine Estuary	49.42	0.25	Upper Galveston Bay Channel	29.68	-94.98
Seto Inland Sea	33.72	132.45	Upper Laguna Madre	27.26	-97.39
Shellbank Basin	40.59	-73.92	Upper Pond, Elkhorn Slough	36.81	-121.74
Shem Creek	32.79	-79.88	Upper Suir Estuary	52.27	-7.0
Singlefjord	59.08	11.12	Vejde Fjord	55.7	9.68
Skagerrak coast (Denmark)	57.38	9.72	Venice Lagoon	45.3	12.4
Skagerrak coast (Fjords)	58.72	10.85	Volga River Delta	45.7	47.9
Skagerrak coast (Sweden)	58.25	11.5	Wadden Sea	53.82	7.67
Skive Fjord	56.62	9.08	Waquoit Bay	41.6	-70.5
Slaney Estuary Lower	52.34	-6.45	Watsons Bayou	30.15	-85.63
Søndeledfjord	58.73	9.15	Wax Lake	29.6	-91.42
South Puget Sound	47.22	-122.81	Weeks Bay	29.79	-84.88
South San Francisco Bay	37.56	-122.18	Wells Inlet	43.35	-70.55
South Slough	43.32	-124.32	Weser Estuary	53.33	8.57
St. Andrew Bay	30.16	-85.7	West Hackberry	29.67	-93.5
St. Anna Archipelago	58.41	16.75	West Mississippi Sound	30.26	-89.02
St. George Sound	29.8	-84.69	Wester-Ems	53.5	6.75
St. Helena Sound	32.54	-80.39	Western Indian Shelf	15.5	73.7
St. Johns River	30.5	-81.5	Westerschelde Estuary	51.4	3.55
St. Joseph Bay	29.79	-85.36	Whidbey Basin & Skagit Bay	48.18	-122.45
St. Lawrence Estuary	48.7	-68.65	Winyah Bay	33.39	-79.26
St. Leonard Creek	38.42	-76.49	Wismar Bay	53.9	11.4
St. Lucie River	27	-80.3	Wolf Bay	30.33	-87.6
Steindalsfjord	58.2	8.32	York River	37.48	-76.75
Stockholm Inner Archipelago	59.47	18.78	Youngsan Estuary	34.83	126.4
Stølefjord	58.82	9.43	Ythan Estuary	57.34	-1.96
Stono & North Edisto Rivers	32.64	-80.03			
Strangford Lough Catchment	54.35	-5.67			
Sullom Voe	60.4	-1.4			
Suo-Nada	33.85	131.17			
Swedish West Coast Fjords	57.92	11.58			
Tampa Bay near Beacon Key	27.74	-82.51			
Tan Shui Estuary	25.17	121.42			
Taskinas Creek	37.41	-76.71			
Taw Estuary	51.08	-4.22			
Tawe Estuary	51.62	-3.93			
Tees Estuary	54.63	-1.15			
Ten Thousand Islands North	25.72	-81.38			
Ten Thousand Islands South	25.31	-81.06			
Thames Estuary	51.8	0.9			
Thau Lagoon	43.67	3.6			

Supplemental References

- Abadie, S.W., M.A. Poirrier. 2000. Increased density of large *Rangia* clams in Lake Pontchartrain after the cessation of shell dredging. *Journal of Shellfish Research* 19:481-485.
- Abril, G., H. Etcheber, P.L. Hir, P. Bassoullet, B. Boutier, M. Frankignoulle. 1999. Oxic/anoxic oscillations and organic carbon mineralization in an estuarine maximum turbidity zone (The Gironde, France). *Limnology and Oceanography* 44:1304-1315.
- Abril, G, H. Etcheber, B. Delille, M. Frankignoulle, A.V. Borges. 2003. Carbonate dissolution in the turbid and eutrophic Loire estuary. *Marine Ecology Progress Series* 259:129-138
- Andersin, A.B., J. Lassig, L. Parkkonen, H. Sandler. 1978. The decline of macrofauna in the deeper parts of the Baltic proper and the Gulf of Finland. *Kieler Meeresforschungen, Sonderheft* 4:23-52.
- Andersin, A.-B., H. Sandler. 1991. Macrobenthic fauna and oxygen deficiency in the Gulf of Finland. *Memoranda Soc Fauna Flora Fennica* 67:3-10.
- Anderson, T.H., G.T. Taylor. 2001. Nutrient pulses, plankton blooms, and seasonal hypoxia in western Long Island Sound. *Estuaries* 24:228-243.
- Andrews, M.J., D.G. Rickard. 1980. Rehabilitation of the inner Thames estuary. *Marine Pollution Bulletin* 11:327-332.
- Araújo F.G., R.G. Bailey, W.P. Williams. 1999. Spatial and temporal variations in fish populations in the upper Thames estuary. *Journal of Fish Biology* 55:836-853
- Arntz, W. 1981. Zonation and dynamics of macrobenthos biomass in an area stressed by oxygen deficiency. *In: Stress effects on natural ecosystems*, Barrett, G.W: and Rosenberg, R. (eds.), John Wiley and Sons, Chichester. pp. 215-225.
- Arvanitidis C., D. Koutsoubas, C. Dounas, A. Eleftheriou. 1999. Annelid fauna of a Mediterranean Lagoon (Gialova Lagoon, south-west Greece): community structure in a severely fluctuating environment. *Journal of the Marine Biological Association, UK* 1999:849-856.
- Baden, S.P., L.O. Loo, L. Pihl, and R. Rosenberg. 1990a. Effects of Eutrophication on benthic communities including fish - Swedish west coast. *Ambio* 19:113-122.
- Baden, S.P., L. Pihl, and R. Rosenberg. 1990b. Effects of oxygen depletion on the ecology, blood physiology and fishery of the Norway lobster *Nephrops norvegicus*. *Marine Ecology Progress Series* 67:141-155.
- Balls P.W., N. Brockie, J. Dobson, W. Johnston. 1996. Dissolved oxygen and nitrification in the upper Forth Estuary during summer (1982-92): patterns and trends. *Estuarine, Coastal and Shelf Science* 42:117-134
- Bakan, G., H. Buyukgungor. 2000. The Black Sea. *Marine Pollution Bulletin* 41:24-43.
- Balkas, T., G. Dechev, R. Mihnea, O. Serbanescu, U. Unlueata. 1990. State of the marine environment in the Black Sea region. *UNEP Regional Seas Reports Studies No. 124*. 41 p.
- Balkis, N. 2003. The effect of Marmara (Iizmit) earthquake on the chemical oceanography of Iizmit Bay, Turkey. *Marine Pollution Bulletin* 46:865-878.
- Barmawidjaja, D.M., G.J. Van Der Zwaan, F.J. Jorissen, S. Puskaric. 1995. 150 years of eutrophication in the northern Adriatic Sea: evidence from a benthic foraminifer record. *Marine Geology* 122:367-384.

- Bentley, S.J., C.A. Nittrouer. 1999. Physical and biological influences on the formation of sedimentary fabric in an oxygen-restricted depositional environment: Eckernförde Bay, Southwestern Baltic Sea. *Palaios* 14.
- Benoit P., Y. Gratton, A. Mucci. 2006. Modeling of dissolved oxygen levels in the bottom waters of the Lower St. Lawrence Estuary: coupling of benthic and pelagic processes. *Marine Chemistry* 102:13-32.
- Bergondo, D.L., D.R. Kester, H.E. Stoffel, W.L. Woods. 2005. Time-series observations during the low sub-surface oxygen events in Narragansett Bay during summer 2001. *Marine Chemistry* 97:90-103.
- Blay, J., Jr., F. Dongdem. 1996. Preliminary observations on the benthic macrofauna of a polluted coastal lagoon in Ghana (West Africa). *Tropical Ecology* 37:127-133.
- Boesch D.F., R.B. Brinsfield, R.E. Magnien. 2001. Chesapeake Bay eutrophication: scientific understanding, ecosystem restoration, and challenges for agriculture. *Journal of Environmental Quality* 30:303-320.
- Bock M.T., B.S. Miller, A.W. Bowman. 1999. Assessment of eutrophication in the Firth of Clyde: analysis of coastal water data from 1982 to 1996. *Marine Pollution Bulletin* 38:222-231.
- Breitburg, D.L, T. Loher, C.A. Pacey, A. Gerstein. 1997. Varying effects of low dissolved oxygen on trophic interactions in an estuarine food web. *Ecological Monographs* 67:489-507.
- Bricker, S.B., C.G. Clement, D.E. Pirhalla, S.P. Orlando, D.R.G. Farrow. 1999. National Estuarine Eutrophication Assessment: Effects of Nutrient Enrichment in the Nation's Estuaries. NOAA, National Ocean Service, Special Projects Office and the National Centers for Coastal Ocean Science. Silver Spring, MD. 71 p.
- Bricker, S., B. Longstaff, W. Dennison, A. Jones, K. Boicourt, C. Wicks, J. Woerner. 2007. Effects of Nutrient Enrichment In the Nation's Estuaries: A Decade of Change. NOAA Coastal Ocean Program Decision Analysis Series No. 26. National Centers for Coastal Ocean Science, Silver Spring, MD. 328 p.
- Brockmann, U., G. Billen, W. Gieskes. 1988. North Sea Nutrients and Eutrophication . *In: Pollution of the North Sea: An Assessment*. W. Salomons, B. Bayne, E. Duursma, U. Förstner (eds.), Springer-Verlag, New York. pp. 348-389.
- Brongersma-Sanders, M. 1957. Mass mortality in the sea. *In: Treatise on Marine Ecology and Paleocology*, Volume 1, J.W. Hedgpeth (ed.), Waverly Press, Baltimore. pp. 941-1010.
- Brosnan, T.M., M.L. O'Shea. 1996. Long-term improvements in water quality due to sewage abatement in the lower Hudson River. *Estuaries* 19:890-900.
- Bukharitsin, P.I., Luneva, Z.D. 1994. Water quality characteristics of the lower Volga reaches and the northern Caspian Sea. *Water Resources* 21:410-416.
- Burnett L.E. 1997. The challenges of living in hypoxic and hypercapnic aquatic environments. *American Zoologist* 37:633-640.
- Chamberlain, R., D. Hayward. 1996. Evaluation of water quality and monitoring in the St. Lucie Estuary, Florida. *Water Resources Bulletin* 32:681-696.
- Chapelle, A., A. Menesguen, J.-M. Deslous-Paoli., P. Souchu, N. Mazouni, A. Vaquer, B. Millet. 2000. Modelling nitrogen, primary production and oxygen in a Mediterranean lagoon. Impact of oysters farming and inputs from the watershed. *Ecological Modelling* 127:161-181.
- Chechun, T.Y. 1998. Nutrition of the Acipenseridae of the Sea of Azov under present conditions. *Journal of Ichthyology* 38:147-154.

- Chen C.-C., G.-C. Gong, F.-K. Shiah. 2007. Hypoxia in the East China Sea: one of the largest coastal low-oxygen areas in the world. *Marine Environmental Research* 64:399-408.
- Cheng W., C.-H. Liu, J.-P. Hsu, J.-C. Chen. 2002. Effect of hypoxia on the immune response of giant freshwater prawn *Macrobrachium rosenbergii* and its susceptibility to pathogen *Enterococcus*. *Fish and Shellfish Immunology* 13:351-365.
- Christensen, J., T. Packard. 1976. Oxygen utilization and plankton metabolism in a Washington fjord. *Estuarine and Coastal Marine Science* 4:339-347.
- Clarke A., S. Juggins, D. Conley. 2003. A 150-year reconstruction of the history of coastal eutrophication in Roskilde Fjord, Denmark. *Marine Pollution Bulletin* 46:1615-1629.
- Cloern J.E., R.S. Oremland. 1983. Chemistry and microbiology of a sewage spill in south San Francisco Bay. *Estuaries* 6:399-406.
- Conley D.J., J. Carstensen, G. Aertebjerg, P.B. Christensen, T. Dalsgaard, J.L.S. Hansen, A.B. Josefson. 2007. Long-term changes and impacts of hypoxia in Danish coastal waters. *Ecological Applications* 17:S165-S184.
- Dai M., X. Guo, W. Zhai, L. Yuan, B. Wang, L. Wang, P. Cai, T. Tang, W.J. Cai. 2006. Oxygen depletion in the upper reach of the Pearl River estuary during a winter drought. *Marine Chemistry* 102:159-169.
- D'Andrea, A.F., N.I. Craig, G.R. Lopez. 1996. Benthic macrofauna and depth of bioturbation in Eckernförde Bay, southwestern Baltic Sea. *Geo-Marine* 16:155-159.
- D'Avanzo C., J.N. Kremer. 1994. Diel oxygen dynamics and anoxic events in an eutrophic estuary of Waquoit Bay, Massachusetts. *Estuaries* 17:131-139.
- Danulat, E., P. Muniz, J. Garcia-Alonso, B. Yannicelli. 2002. First assessment of the highly contaminated harbour of Montevideo, Uruguay. *Marine Pollution Bulletin* 44:554-565.
- Dauer D.M., A.J. Rodi, Jr., J.A. Ranasinghe. 1992. Effects of low dissolved oxygen events on the macrobenthos of the lower Chesapeake Bay. *Estuaries* 15:384-391.
- Deacutis, C.F., D. Murray, W. Prell, E. Saarman, L. Korhun. 2006. Hypoxia in the upper half of Narragansett Bay, RI, during August 2001 and 2002. *Northeastern Naturalist* 13:173-198.
- Debolskaya E.I., E.V. Yakushev, I.S. Kuznetsov. 2008. Analysis of the hydrophysical structure of the Sea of Azov in the period of the bottom anoxia development. *Journal of Marine Science* 70:300-307.
- DeJonge V., W. Boynton, C. D'Elica, R. Elmgren, R. Welsh. 1994. Responses to developments in eutrophication in four different North Atlantic estuarine systems. *In: Changes in fluxes in estuaries: implications from science to management*. K. Dyer, and R. Orth (eds.), Olsen and Olsen, Fredensborg.
- Desprez, M., H. Rybarczyk, J.G. Wilson, J.P. Ducrottoy, F. Sueur, R. Olivesi, B. Elkaim. 1992. Biological impact of eutrophication in the Bay of Somme and the induction and impact of anoxia. *Netherlands Journal of Sea Research* 30:149-159.
- Dethlefsen, V., H. v. Westernhagen. 1983. Oxygen deficiency and effects on bottom fauna in the eastern German Bight 1982. *Meeresforschung* 60:42-53.
- Diaz, R.J., R.J. Neubauer, L.C. Schaffner, L. Pihl, S.P. Baden. 1992. Continuous monitoring of dissolved oxygen in an estuary experiencing periodic hypoxia and the effect of hypoxia on macrobenthos and fish. *Science of the Total Environment, Supplement* 1992:1055-1068.

- Dommasnes, A., F. Rey, I. Rottingen. 1994. Reduced oxygen concentration in herring wintering areas. *ICES Journal of Marine Science* 51:63–69.
- Donnelly, T.H., P.W. Ford, D. McGregor, D. Allen. 1999. Anthropogenic changes to a billabong in New South Wales. 1. Lagoon evolution and phosphorus dynamics. *Marine and Freshwater Research* 50:689-698.
- Dorgham, M.M., N.E. Abdel-Aziz, K.Z. ElDeeb, M.A. Okbah. 2004. Eutrophication problem in the Western Harbour of Alexandria, Egypt. *Oceanologia* 46:1-20.
- Doulgeraki, S., N. Lampadariou, A. Sinis. 2006. Meiofaunal community structure in three Mediterranean coastal lagoons (North Aegean Sea). *Journal of the Marine Biological Association of the United Kingdom* 86:209-220.
- Dumont, H.J. 1998. The Caspian Lake: History, biota, structure, and function. *Limnology and Oceanography* 43:44-52.
- Ducrotoy J.-P., M. Elliott, V.N.D. Jonge. 2000. The North Sea. *Marine Pollution Bulletin* 41:5-23.
- Dyer, M.F., J.G. Pope, P.D. Fry, R.J. Law, J.E. Portmann. 1983. Changes in fish and benthos catches off the Danish coast in September 1981. *Journal of the Marine Biological Association of the United Kingdom* 63:767-775.
- Eden N., T. Katz, D.L. Angel. 2003. Dynamic response of a mud snail *Nassarius sinusigerus* to changes in sediment biogeochemistry. *Marine Ecology Progress Series* 263:139-147.
- Engle, V.D., J.K. Summers, J.M. Macauley. 1999. Dissolved oxygen conditions in northern Gulf of Mexico estuaries. *Environmental Monitoring and Assessment* 57:1-20.
- Engle, V.D., J.K. Summers. 1999. Refinement, validation, and application of a benthic condition index for northern Gulf of Mexico estuaries. *Estuaries* 22:624-635.
- Erez, J., D. Iluz, D. Zakai, J. Silverman, B. Lazar. 2004. Eutrophication processes in the Gulf of Eilat (Aqaba), Red-Sea, and their effects on the coral reef ecosystem. <http://adsabs.harvard.edu/abs/2003EAEJA....11772E>
- Fallesen, G., H.M. Jørgensen. 1991. Distribution of *Nephtys hombergii* and *N. ciliata* (Polychaeta: Nephtyidae) in Århus Bay, Denmark, with emphasis on the effect of sewage oxygen deficiency. *Ophelia Supplement* 5:443-450.
- Fallesen, G., F. Andersen, B. Larsen. 2000. Life, death and revival of the hypertrophic Mariager Fjord, Denmark. *Journal of Marine Systems* 25:313-321.
- Fedra K., E.M. Ölscher, C. Scherübel, M. Stachowitsh, R.S. Wurzian. 1976. On the ecology of a North Adriatic benthic community: distribution, standing crop and composition of the macrobenthos. *Marine Biology* 38:129-145.
- Flemer, D.A., W.L. Kruczynski, B.F. Ruth, C.M. Bundrick. 1999. The relative influence of hypoxia, anoxia, and associated environmental factors as determinants of macrobenthic community structure in a northern Gulf of Mexico. *Journal of Aquatic Ecosystem Stress and Recovery* 6:311-328.
- Flindt M.R., L. Kamp-Nielsen, J.C. Marques, M.A. Pardal, M. Bocci, G. Bendoricchio, J. Salomonsen, S.N. Nielsen, S.E. Jørgensen. 1997. Description of the three shallow estuaries: Mondego River (Portugal), Roskilde Fjord (Denmark) and the Lagoon of Venice (Italy). *Ecological Modeling* 102:17-31.
- Fonseca, A. 2006. Efeito da drenagem urbana nas características físico-químicas e biológicas da água superficial na Lagoa da Conceição (Florianópolis, SC, Brasil). *Biotemas* 19:7-16.
- Foulkes, M., S. Rattanachongkiat, W. Utoomprukporn, M. Taiyaqun, P. Tantichodok, P. Chongprasith, G. Millward.

2007. Water chemistry and arsenic concentrations in Pak Panang Bay, southern Thailand: influences of the north east monsoon. *Journal of Coastal Research* 23:731-739.
- Friedrich, J., C. Dinkel, G. Friedl, N. Pimenov, J. Wijsman, M.-T. Gomoiu, A. Cociasu, L. Popa, B. Wehrli. 2002. Benthic nutrient cycling and diagenetic pathways in the north-western Black Sea. *Estuarine, Coastal and Shelf Science* 54:369–383.
- Friligos, N., A. Zenetos. 1988. Elefsis Bay anoxia: nutrient conditions and benthic community structure. *Marine Ecology* 9:273-290.
- Fritz, C., E. LaBrecque, J. Tober, P.J. Behr, I. Valiela. 1996. Shrimp in Waquoit Bay: Effects of nitrogen loading on size and abundance. General Scientific Meetings of the Marine Biological Laboratory, Woods Hole, MA 191:326-327.
- Garlo, E.V., C.B. Milstein, A.E. Jahn. 1979. Impact of hypoxic conditions in the vicinity of Little Egg Inlet, New Jersey in summer 1976. *Estuarine and Coastal Marine Science* 8:421-432.
- Garnier J., P. Servais, G. Billen, M. Akopian, N. Brion. 2001. The Seine River and Estuary (France) carbon and oxygen budgets during low flow. *Estuaries* 24:964-976.
- Gage J. 1972. A preliminary survey of the benthic macrofauna and sediments in lochs Etive and Creran, sea-lochs along the west coast of Scotland. *Journal of the Marine Biological Association, UK* 52:237-276.
- Gaston G. 1985. Effects of hypoxia on macrobenthos of the inner shelf off Cameron, Louisiana. *Estuarine, Coastal and Shelf Science* 20:603-613.
- Gerlach, S.A. 1990. Nitrogen, phosphorus, plankton and oxygen deficiency in the German Bight and in Kiel Bay. Final Report, Eutrophication of the North Sea and the Baltic Sea. *Kieler Meeresforschungen, Sonderheft Nr. 7.* 332 p.
- Gibbs, P.E., J.C. Green, P.L. Pascoe. 1999. A massive summer-kill of the dog whelk, *Nucella lapillus*, on the north Cornwall coast in 1995: freak or forerunner? *Journal of the Marine Biological Association, United Kingdom* 79:103-109.
- Gilbert, D., B. Sundby, C. Gobeil, A. Mucci, G.-H. Tremblay. 2005. A seventy-two-year record of diminishing deep-water oxygen in the St. Lawrence estuary: The northwest Atlantic connection. *Limnology and Oceanography* 50:1654-1666.
- Gillett D.J., A.F. Holland, D.M. Sanger. 2005. Secondary production of a dominant oligochaete (*Monopylephorus rubroniveus*) in the tidal creeks of South Carolina and its relation to ecosystem characteristics. *Limnology and Oceanography* 50:566-577.
- Gillibrand, P.A., W.R. Turrell, D.C. Moore, R.D. Adams. 1996. Bottom water stagnation and oxygen depletion in a Scottish sea loch. *Estuarine, Coastal and Shelf Science* 43:217-235.
- Glenn, S., M. Crowley, D. Haidvogel, Y. Song. 1996. Underwater observatory captures coastal upwelling event off New Jersey. *Eos, Transactions of the American Geophysical Union* 77:223-236.
- Gonzales-Oreja, J., J. Saiz-Salinas. 1998. Loss of heterotrophic biomass structure in an extreme estuarine environment. *Estuarine, Coastal and Shelf Science* 48:391-399.
- Grantham B.A., F. Chan, K.J. Nielsen, D.S. Fox, J.A. Barth, A. Huyer, J. Lubchenco, B.A. Menge. 2004. Upwelling-driven nearshore hypoxia signals ecosystem and oceanographic changes in the northeast Pacific. *Nature* 429:749-754.
- Griffiths, A.H. 1987. Water quality of the estuary and Firth of Forth, Scotland. *Proceedings of the Royal Society of*

- Edinburgh 93B:303–314.
- Gunter, G. 1942. Offatts Bayou, a locality with recurrent summer mortality of marine organisms. *American Midland Naturalist* 28:631-633.
- Gustafsson M., K. Nordberg 1999. Benthic foraminifera and their response to hydrography, periodic hypoxic conditions and primary production in the Koljo fjord on the Swedish west coast. *Journal of Sea Research* 41:163-178.
- Gustafsson M., K. Nordberg. 2000. Living (stained) benthic foraminifera and their response to the seasonal hydrographic cycle, periodic hypoxia and to primary production in Havstens Fjord on the Swedish west coast. *Estuarine, Coastal and Shelf Science* 51:743-761.
- Gustafsson M., K. Nordberg. 2001. Living (stained) benthic foraminiferal response to primary production and hydrography in the deepest part of the Gullmar Fjord, Swedish west coast, with comparisons to Høglund's 1927 material. *Journal of Foraminiferal Research* 31:2-11.
- Guyoneaud, R., R. De Wit, R. Matheron, P. Caumette. 1998. Impact of macroalgal dredging on dystrophic crises and phototrophic bacterial blooms (red waters) in a brackish coastal lagoon. *Oceanologica Acta* 21:551-561.
- Hagy, J.D., W.R. Boynton, C.W. Keefe, K.V. Wood. 2004. Hypoxia in Chesapeake Bay, 1950-2001: Long-term change in relation to nutrient loading and river flow. *Estuaries* 27:634-658.
- Harper D.E., Jr., N.N. Rabalais. 1995. Responses of Benthonic and Nektonic Organisms, and communities, to severe hypoxia on the inner continental shelf of Louisiana and Texas. *In: Proceedings of the First Gulf of Mexico Hypoxia Management Conference, Dec 5-6 1995, Kenner, Louisiana, USEPA-55-R-97-002.* pp. 41-56.
- Harper, D.E., L.D. McKinney, J.M. Nance, R.B. Salzer. 1991. Recovery responses of two benthic assemblages following an acute hypoxic event on the Texas continental shelf, northwestern Gulf of Mexico. *In: Modern and ancient continental shelf anoxia.* R.V. Tyson and T.H. Pearson (eds.). Geological Society Special Publication No. 58, London. pp. 49-64.
- Harper, D.E., L.D. McKinney, R.B. Salzer, R.J. Case. 1981. The occurrence of hypoxic bottom water off the upper Texas coast and its effects on the benthic biota. *Contributions in Marine Science* 24:53-79.
- Hearn C.J., B.J. Robson. 2001. Inter-annual Variability of Bottom Hypoxia in Shallow Mediterranean Estuaries. *Estuarine, Coastal and Shelf Science* 52:643-657.
- Hendelberg, M., P. Jensen. 1993. Vertical distribution of the nematode fauna in a coastal sediment influenced by seasonal hypoxia in the bottom water. *Ophelia* 37:83-94.
- Hobbie, J.E., B.J. Copeland, W.G. Harrison. 1975. Sources and fates of nutrients in the Pamlico River estuary of North Carolina. *In: L. E. Cronin (ed.), Estuarine Research. Vol. 1.* Academic Press, New York. pp. 287-302.
- Hodoki Y., T. Murakami. 2006. Effects of tidal flat reclamation on sediment quality and hypoxia in Isahaya Bay. *Aquatic Conservation-Marine and Freshwater Ecosystems* 16:555-567.
- Holland, A.F., A.T. Shaughnessy, M.H. Hiegel. 1987. Long-term variation in mesohaline Chesapeake Bay macrobenthos: spatial and temporal patterns. *Estuaries* 10:370-378.
- Howell, P., D. Simpson. 1994. Abundance of marine resources in relation to dissolved oxygen in Long Island Sound. *Estuaries* 17:394-402.

- Hylleberg, J. 1993. Extinction and immigration of benthic fauna, the value of historical data from Limfjorden, Denmark. Symposium Mediterranean Seas 2000, Santa Margherita Ligure, 23-27. September 1991, Ist. Sci. Ambientali Mar., Santa Margherita Ligure, Italy. pp. 43-70.
- Iizuka, S., S. Min. 1989. Formation of anoxic bottom waters in Omura Bay. *Engan Kaiyo Kenkyu* 2:75-86.
- Imabayashi, H. 1986. Effect of oxygen-deficient water on the settled abundance and size composition of the bivalve *Theora lubrica*. *Bulletin of the Japanese Society of Scientific Fisheries* 5:391-397.
- Irigoién X., J. Post, J. Castel, K.F. Pfeiffer, B. Hellmann. 1999. Nycthemeral variations of the dissolved oxygen concentration in the turbidity maximum of three European estuaries: biological vs. physical processes. *Journal of Marine Systems* 22:173-177.
- Isaza, C.F.A., P.C. Sierra-Correa, M. Bernal-Velasquez, L.M. Londoño, W. Troncoso. 2006. Caribbean Sea/Colombia & Venezuela, Caribbean Sea/Central America & Mexico, GIWA Regional assessment 3b, 3c. UNEP. University of Kalmar, Kalmar, Sweden. UNEP, 2006.
- Jeng, W.-L., B.-C. Han. 1996. Coprostanol in a sediment core from the anoxic Tan-Shui Estuary, Taiwan. *Estuarine, Coastal and Shelf Science* 42:727-735.
- Johannessen T., E. Dahl. 1996a. Declines in oxygen concentrations along the Norwegian Skagerrak coast, 1927-1993: A signal of ecosystem changes due to eutrophication? *Limnology and Oceanography* 41:766-778.
- Johannessen T., E. Dahl. 1996b. Historical changes in oxygen concentrations along the Norwegian Skagerrak coast: Reply to the comment by Gray and Abdullah. *Limnology and Oceanography* 41:1847-1852.
- Johansson, J.O.R., R.R. Lewis III. 1992. Recent improvements of water quality and biological indicators in Hillsborough Bay, a highly impacted subdivision of Tampa Bay, Florida, USA. *In*: Vollenweider, R.W., R. Marchetti and R. Viviani (eds.), *Marine coastal eutrophication: The response of marine transitional systems to human impact: Problems and perspectives for restoration*. New York. Elsevier Press. pp. 1191-1215.
- Jones P.D. 2006. Water quality and fisheries in the Mersey estuary, England: A historical perspective. *Marine Pollution Bulletin* 53:144-154
- Jørgensen, B.B. 1980. Seasonal oxygen depletion in the bottom waters of a Danish fjord and its effect on the benthic community. *Oikos* 34:68-76.
- Josefson A.B., J.L.S. Hansen. 2004. Species richness of benthic macrofauna in Danish estuaries and coastal areas. *Global Ecology and Biogeography* 13:273-288
- Josefson, A.B., J.N. Jensen. 1992. Effects of hypoxia on soft-sediment macrobenthos in southern Kattegat. *In*: *Marine eutrophication and population dynamics*. G. Colombo, I. Ferrari, V. U. Ceccherelli and R. Rossi (eds.), Olsen and Olsen, Fredensborg. pp. 21-28.
- Josefson, A.B., R. Rosenberg. 1988. Long-term soft-bottom faunal changes in three shallow fjords, west Sweden. *Netherlands Journal of Sea Research* 22, 149-159.
- Josefson, A.B., B. Widbom. 1988. Differential response of benthic macrofauna and meiofauna to hypoxia in the Gullmar Fjord basin. *Marine Biology* 100:31-40.
- Justic', D., T. Legovic', L. Rottini-Sandrini. 1987. Trends in oxygen content 1911-1984 and occurrence of benthic mortality in the northern Adriatic Sea. *Estuarine Coastal Shelf Science* 25:435-445.
- Justic', D., N.N. Rabalais, R.E. Turner. 1996. Effects of climate change on hypoxia in coastal waters: a doubled CO₂ scenario for the northern Gulf of Mexico. *Limnology and Oceanography* 41:992-1003.

- Justic', D., N.N. Rabalais, R.E. Turner, W.J. Wiseman, Jr. 1993. Seasonal coupling between riverborne nutrients, net productivity and hypoxia. *Marine Pollution Bulletin* 26:184-189.
- Kahru, M., B.G. Mitchell, A. Diaz, M. Miura. 2004. MODIS detects a devastating algal bloom in Paracas Bay, Peru. *Eos, Transactions of the American Geophysical Union* 85:465-472.
- Kaiser, J., S. Lutter. 1998. Do we have the right strategies to combat eutrophication in the Wadden Sea? A critical review of current policies. *Ecosystem Research in the Wadden Sea Area*. 9. International Scientific Wadden Sea Symposium, Norderney, Germany, 5-8 Nov 1996. *Senckenbergiana maritima* 29:17-24.
- Karim, M.R., S. Masahiko, U. Masao. 2002. Simulation of eutrophication and associated occurrence of hypoxic and anoxic condition in a coastal bay in Japan. *Marine Pollution Bulletin* 45:280-285.
- Karlson, K., R. Rosenberg, E. Bonsdorff. 2002. Temporal and spatial large-scale effects of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic waters – a review. *Oceanography and Marine Biology: an Annual Review* 40:427-489.
- Kasai, A., H. Toyohara, A. Nakata, T. Miura, N. Azuma. 2006. Food sources for the bivalve *Corbicula japonica* in the foremost fishing lakes estimated from stable isotope analysis. *Fisheries Science* 72:105-114.
- Keister, J.E., E.D. Houde, D.L. Breitburg. 2000. Effects of bottom-layer hypoxia on abundances and depth distributions of organisms in Patuxent River, Chesapeake Bay. *Marine Ecology Progress Series* 205:43-59.
- Kelly, M., M. Naguib. 1984. Eutrophication in coastal marine areas and lagoons: a case study of 'Lac de Tunis'. *UNESCO Reports in Marine Science*. Vol. 29. 71 p.
- Kim, D.-H., O. Matsuda, T. Yamamoto. 1997. Nitrification, denitrification and nitrate reduction rates in the sediment of Hiroshima Bay, Japan. *Journal of Oceanography* 53:317-324.
- Kimura M., T. Takahashi, T. Takatsu, T. Nakatani, T. Maeda. 2004. Effects of hypoxia on principal prey and growth of flathead flounder *Hippoglossoides dubius* in Funka Bay, Japan. *Fisheries Science* 70:537-545.
- Kitching, J.A., F.J. Ebling, J.C. Gable, R. Hoare, A.A.Q.R. McLeod, T.A. Norton. 1976. The ecology of Lough Ine. XIX. Seasonal changes in the western trough. *Journal of Animal Ecology* 45:731-758.
- Kodama K., I. Aoki, M. Shimizu. 2002. Long-term changes in the assemblage of demersal fishes and invertebrates in relation to environmental variations in Tokyo Bay, Japan. *Fisheries Management and Ecology* 9:303-313.
- Kodama K., T. Horiguchi, G. Kume, S. Nagayama, T. Shimizu, H. Shiraishi, M. Morita, M. Shimizu. 2006. Effects of hypoxia on early life history of the stomatopod *Oratosquilla oratoria* in a coastal sea. *Marine Ecology Progress Series* 324:197-206.
- Koepfler, E., S. Lake, E.M. Smith, J. Bennett, S. Libes. 2007. Examination of Inner Shelf Water Quality in Long Bay, South Carolina, using Dataflow. *Estuarine Research Federation Providence Rhode Island Meeting*, November 2007, abstract.
- Koutrakis E.T., N.I. Kamidis, I.D. Leonardos. 2004. Age, growth and mortality of a semi-isolated lagoon population of sand smelt, *Atherina boyeri* (Risso, 1810) (Pisces: Atherinidae) in an estuarine system of northern Greece. *Journal of Applied Ichthyology* 20:382-388.
- Kozlowsky-Suzuki, B., R.L. Bozelli. 2002. Experimental evidence of the effect of nutrient enrichment on the zooplankton in a Brazilian coastal lagoon. *Brazilian Journal of Biology* 62:835-846.

- Kraufvelin P., B. Sinisalo, E. Leppakoski, J. Mattila, E. Bonsdorff. 2001. Changes in zoobenthic community structure after pollution abatement from fish farms in the Archipelago Sea (N. Baltic Sea). *Marine Environmental Research* 51:229-245.
- Laine, A.O., H. Sandler, A-B. Andersin, J. Stigzelius. 1997. Long-term changes of macrozoobenthos in the eastern Gotland Basin and the Gulf of Finland (Baltic Sea) in relation to the hydrographical regime. *Netherlands Journal of Sea Research* 38:135-159.
- Lapointe, B.E., W.R. Matzie. 1996. Effects of stormwater nutrient discharges on eutrophication processes in nearshore waters of the Florida Keys. *Estuaries* 19:422-435.
- Lardicci C., S. Como, S. Corti, F. Rossi. 2001. Recovery of the macrozoobenthic community after severe dystrophic crises in a Mediterranean coastal lagoon (Orbetello, Italy). *Marine Pollution Bulletin* 42:202-214.
- Le Bris H., M. Glemarec. 1995. Macrobenthic communities of an oxygen of under-saturated coastal ecosystems: The Bay of Vilaine (southern Brittany). *Oceanologica Acta* 18:573-581.
- Lee, M.-O., J.-K. Kim. 2008. Characteristics of algal blooms in the southern coastal waters of Korea. *Marine Environmental Research* 65:128-147.
- Legovic', T., D. Petricioli, V. Zutic. 1991. Hypoxia in a pristine stratified estuary (Krka, Adriatic Sea). *Marine Chemistry* 32:347-359.
- Lehman P.W., J. Sevier, J. Giulianotti, M. Johnson. 2004. Sources of oxygen demand in the lower San Joaquin River, California. *Estuaries* 27:405-418.
- Le Loeuff, P. 1999. The benthic macrofauna of the variable salinity waters ecosystems along the Atlantic coast of tropical Africa; biodiversity variations with the current climatic conditions (rainfall) and the regional climatic history. *Zoosystema* 21:557-571.
- Lenihan H.S. 1999. Physical-biological coupling on oyster reefs: how habitat structure influences individual performance. *Ecological Monographs* 69:251-275.
- Lenihan, H.S., C.H. Peterson. 1998. How habitat degradation through fishery disturbance enhances impacts of hypoxia on oyster reefs. *Ecological Applications* 8:128-140.
- Leonard, C.L., J.B. McClintock. 1999. The population dynamics of the brittlestar *Ophioderma brevispinum* in near- and farshore seagrass habitats of Port Saint Joseph Bay, Florida. *Gulf of Mexico Science* 17:87-94.
- Leppäkoski, E. 1969. Transitory return of the benthic fauna of the Bornholm Basin, after extermination by oxygen insufficiency. *Cahiers de Biologie Marine* 10:163-172.
- Lerberg S.B., A.F. Holland, D.M. Sanger. 2000. Responses of tidal creek macrobenthic communities to the effects of watershed development. *Estuaries* 23:838-853.
- Leverone, J.R. 1995. Diurnal dissolved oxygen in two Tampa Bay seagrass meadows: ramifications for the survival of adult bay scallops (*Argopectin irradians concentricus*). *Florida Scientist* 58:141-152.
- Li, D., D. Daler. 2004. Ocean pollution from land-based sources: East China Sea, China. *Ambio* 33:107-113.
- Lim H.S., R.J. Diaz, J.S. Hong, L.C. Schaffner. 2006. Hypoxia and benthic community recovery in Korean coastal waters. *Marine Pollution Bulletin* 52:1517-1526.
- Lin H.-J., K.-T. Shao, R.-Q. Jan, H.-L. Hsieh, C.-P. Chen, L.-Y. Hsieh, Y.-T. Hsiao. 2007. A trophic model for the Danshuei River Estuary, a hypoxic estuary in northern Taiwan. *Marine Pollution Bulletin* 54:1789-1800.

- Lindholm, T. 1996. Periodic anoxia in an emerging coastline lake basin in SW Finland. *Hydrobiologia* 325:223-230.
- Llansó, R.J. 1992. Effects of hypoxia on estuarine benthos: the lower Rappahannock River (Chesapeake Bay), a case study. *Estuarine, Coastal and Shelf Science* 35:491-515.
- Loesch H. 1960. Sporadic mass shoreward migrations of demersal fish and crustaceans in Mobile Bay, Alabama. *Ecology* 41:292-98.
- Lopes J.F., J.M. Dias, A.C. Cardoso, C.I.V. Silva. 2005. The water quality of the Ria de Aveiro lagoon, Portugal: From the observations to the implementation of a numerical model. *Marine Environmental Research* 60:594-628.
- Lopez-Jamar, E., O. Francesch, A.V. Dorrio, S. Parra. 1995. Long-term variation of the infaunal benthos of La Coruna Bay (NW Spain): results from a 12-year study (1982-1993). *Scientia Marina* 59:49-61.
- Lowery, T.A. 1998. Modelling estuarine eutrophication in the context of hypoxia, nitrogen loadings, stratification and nutrient ratios. *Journal of Environmental Management* 52:289-305.
- Luther, G.W., S.F. Ma, R. Trouwborst, B. Glazer, M. Blickley, R.W. Scarborough, M.G. Mensinger. 2004. The roles of anoxia, H₂S, and storm events in fish kills of dead-end canals of Delaware inland bays. *Estuaries* 27:551-560.
- Maciolek, N.J., R.J. Diaz, D.T. Dahlen, C.D. Hunt, I.P. Williams. 2005. 2003 Boston Harbor Benthic Monitoring Report. Boston: Massachusetts Water Resources Authority. Report ENQUAD 2005-06. 83 p.
- MacPherson, T.A., L.B. Cahoon, and M.A. Mallin. 2007. Water column oxygen demand and sediment oxygen flux: patterns of oxygen depletion in tidal creeks. *Hydrobiologia* 586:235-248.
- Mallin, M.A., M.H. Posey, G.C. Shank, M.R. McIver, S.H. Ensign, T.D. Alphin. 1999. Hurricane effects on water quality and benthos in the Cape Fear watershed: Natural and anthropogenic impacts. *Ecological Applications* 9:350-362.
- Marques, M., B. Knoppers, A.E. Lanna, P.R. Abdallah, M. Polette. 2004. Brazil Current, GIWA Regional assessment 39. UNEP. University of Kalmar, Kalmar, Sweden. 192 p.
- Mason, W.T., Jr. 1998. Macrobenthic monitoring in the Lower St. Johns River, Florida. *Environmental Monitoring and Assessment* 50:101-130.
- Maxted J.R., R.A. Eskin, S.B. Weisberg, J.C. Chaillou, F.W. Kutz. 1997. The ecological condition of dead-end canals of the Delaware and Maryland coastal bays. *Estuaries* 20:319-327.
- May, E. 1973. Extensive oxygen depletion in Mobile Bay, Alabama. *Limnology and Oceanography* 18:353-366.
- Mazouni, N., J.-C. Gaertner, J.M. Deslous-Paoli, S. Landrein, M. Geringer d'Oedenberg. 1996. Nutrient and oxygen exchanges at the water-sediment interface in a shellfish farming lagoon (Thau, France). *Journal of Experimental Marine Biology and Ecology* 205:91-113.
- Mee, L. 2006. Reviving dead zones. *Scientific American* 295:78-85.
- Miron, G., T. Landry, P. Archambault, B. Frenette. 2005. Effects of mussel culture husbandry practices on various benthic characteristics. *Aquaculture* 250:138-154.
- Mirza, F.B., J.S. Gray. 1981. The fauna of benthic sediments from the organically enriched Oslofjord, Norway. *Journal of Experimental Marine Biology and Ecology* 54:181-207.

- Montagna, P.A., R.D. Kalke. 1992. The effect of freshwater inflow on meiofaunal and macrofaunal populations in the Guadalupe and Nueces Estuaries, Texas. *Estuaries* 15:307-326.
- Montagna, P.A., C. Ritter. 2006. Direct and indirect effects of hypoxia on benthos in Corpus Christi Bay, Texas, USA. *Journal of Experimental Marine Biology and Ecology* 330:119-131.
- Morrisey, D.J. 2000. Predicting impacts and recovery of marine-farm sites in Stewart Island, New Zealand, from the Findlay-Watling model. *Aquaculture* 185:257-271.
- Moser, F.C. 1998. Sources and sinks of nitrogen and trace metals, and benthic macrofauna assemblages in Barnegat Bay, New Jersey. Dissertation Abstracts International Part B: Science and Engineering, University Microfilms International 58:5849.
- Nagy, G.J., M. Go´mez-Erache, C.H. Lo´pez, A.C. Perdomo. 2002. Distribution patterns of nutrients and symptoms of eutrophication in the Rio de la Plata River Estuary System. *Hydrobiologia* 475/476:125-139.
- Nakata, K., S-I. Sugioka, T. Hosaka. 1997. Hindcast of a Japan Sea Oil Spill. *Spill Science and Technology Bulletin* 4:219-229.
- Naqvi, S.W.A., D.A. Jayakumar, P.V. Narvekar, H. Naik, V. Sarma, W. D'Souza, S. Joseph, M.D. George. 2000. Increased marine production of N₂O due to intensifying anoxia on the Indian continental shelf. *Nature* 408:346-349.
- Neponset River Watershed Association (NERWA). 2004. Boston Harbor south watersheds 2004 assessment report. Massachusetts Executive Office of Environmental Affairs, Boston. 180 p.
- Newcombe, C.L., W.A. Horne. 1938. Oxygen-poor waters of the Chesapeake Bay. *Science* 88:80-81.
- Nichols, F.H., J.E. Cloern, S.N. Luoma, D.H. Peterson. 1986. The modification of an estuary. *Science* 231:567-573.
- Niermann, U., E. Bauerfeind, W. Hickel, H.V. Westernhagen. 1990. The recovery of benthos following the impact of low oxygen content in the German Bight. *Netherlands Journal of Sea Research* 25:215-226.
- Nilsson H., R. Rosenberg. 2000. Succession in marine benthic habitats and fauna in response to oxygen deficiency: analysed by sediment profile-imaging and by grab samples. *Marine Ecology Progress Series* 197:139-149.
- Nobre, A.M., J.G. Ferreira, A. Newton, T. Simas, J.D. Icelly, R. Neves. 2005. Management of coastal eutrophication: Integration of field data, ecosystem-scale simulations and screening models. *Journal of Marine Systems* 56:375-390.
- Norkko, A., E. Bonsdorff. 1996. Rapid zoobenthic community responses to accumulations of drifting algae. *Marine Ecology Progress Series* 131:143-157.
- Oeschger, A., K.B. Storey. 1990. Regulation of glycolytic enzymes in the marine invertebrate *Halicryptus spinulosus* (Priapulida) during environmental anoxia and exposure to hydrogen sulfide. *Marine Biology* 106:261-266.
- Officer, C.B., R.B. Biggs, J.L. Taft, L.E. Cronin, M.A. Tyler, W.R. Boynton. 1984. Chesapeake Bay anoxia: origin, development, and significance. *Science* 223:22-27.
- Ojaveera, H., A. Andrushaitis. 2004. History of ecosystem studies of the Gulf of Riga (Baltic Sea). *Proceedings of the Estonian Academy of Science Biology and Ecology* 53:116-143.
- Okey, T.A. 2003. Macrobenthic colonist guilds and renegades in Monterey canyon (USA) drift algae: partitioning multidimensions. *Ecological Monographs* 73:415-440.

- OSPAR Commission. 2003. OSPAR integrated report 2003 on the eutrophication status of the OSPAR maritime area based upon the first application of the comprehensive procedure, London, U.K. 59 p.
- Paerl, H.W., J.L. Pinkney, S.A. Kucera. 1995. Clarification of the structural and functional roles of heterocysts and anoxic microzones in the control of pelagic nitrogen fixation. *Limnology and Oceanography* 40:634-638.
- Paerl, H.W., J.L. Pinckney, J.M. Fear, B.L. Peierls. 1998. Ecosystem responses to internal and watershed organic matter loading: Consequences for hypoxia in the eutrophying Neuse River Estuary, North Carolina, USA. *Marine Ecology Progress Series* 166:17-25.
- Paerl, H.W., J.D. Bales, L.W. Ausley, C.P. Buzzelli, L.B. Crowder, L.A. Eby, M. Go, B.L. Peierls, T.L. Richardson, J.S. Ramus. 2000. Hurricane's hydrological, ecological effects linger in major US estuary. *Eos, Transactions of the American Geophysical Union* 81:457-462.
- Patrick, R. 1988. Changes in the chemical and biological characteristics of the Upper Delaware River estuary in response to environmental laws. *In: Majumdar, E., W. Miller, and L.E. Sage, Pennsylvania Academy of Science, Philadelphia, PA.* pp. 332-359.
- Parker, C.A., J.E. O'Reilly. 1991. Oxygen depletion in Long Island Sound: A historical perspective. *Estuaries* 14:248-264.
- Paulson, A.J., H.C. Curl, Jr., R.A. Freely. 1992. The biogeochemistry of nutrients and trace metals in Hood Canal, a Puget Sound fjord. *Marine Chemistry* 43:157-173.
- Pearson, T.H., A. Eleftheriou. 1981. The benthic ecology of Sollom Voe. *Proceedings of the Royal Society of Edinburgh* 80B:241-269.
- Pennock, J.R., J.H. Sharp, W.W. Schroeder. 1994. What Controls the Expression of Estuarine Eutrophication? Case Studies of Nutrient Enrichment in the Delaware Bay and Mobile Bay Estuaries, USA. *In: Changes in Fluxes in Estuaries: Implications from Science to Management.* Edited by K.R. Dyer and R.J. Orth. Olsen and Olsen, Fredensborg. p. 139-146.
- Petersen, C.G.J. 1915. On the animal communities of the sea bottom in the Skagerak, the Christiania Fjord and the Danish waters. *Report from the Danish Biological Station* 23:1-28.
- Petersen, J.K., L. Pihl. 1995. Responses to hypoxia of plaice, *Pleuronectes platessa* and dab, *Limanda limanda*, in the south-east Kattegat: distribution and growth. *Environmental Biology of Fishes* 43:311-321.
- Pihl, L., S.P. Baden, R.J. Diaz. 1991. Effects of periodic hypoxia on distribution of demersal fish and crustaceans. *Marine Biology* 108:349-360.
- Pohl, C., U. Hennings, I. Petersohn, H. Siegel. 1998. Trace metal budget, transport, modification and sink in the transition area between the Oder and Peene rivers and the southern Pomeranian Bight. *Marine Pollution Bulletin* 36:598-616.
- Portnoy, J.W. 1991. Summer Oxygen Depletion in a Diked New England Estuary. *Estuaries* 14:122-129.
- Posey, M.H., T.D. Alphin, L. Cahoon, D. Lindquist, M.E. Becker. 1999. Interactive effects of nutrient additions and predation on infaunal communities. *Estuaries* 22:785-792.
- Powilleit, M., J. Kube. 1999. Effects of severe oxygen depletion on macrobenthos in the Pomeranian Bay (southern Baltic Sea): a case study in a shallow, sublittoral habitat characterised by low species richness. *Netherlands Journal of Sea Research* 42:221-234.
- Prena, J. 1995a. Effects of eutrophication on macrobenthos zonation in Wismar Bay (western Baltic Sea). *Archiv fur Hydrobiologie Stuttgart* 133:245-257

- Prena, J. 1995b. Temporal irregularities in the macrobenthic community and deep-water advection in Wismar Bay (western Baltic Sea). *Estuarine, Coastal and Shelf Science* 41:705-717.
- Puente, A., J.A. Juanes, A. Garcí'a, C. A'lvarez, J.A. Revilla, I. Carranza. 2007. Ecological assessment of soft bottom benthic communities in northern Spanish estuaries. *Ecological Indicators* (In Press).
- Rabalais, N.N. 1998. Oxygen depletion in coastal waters. National Oceanic and Atmospheric Administration's State of the Coast Report. Silver Spring, MD: NOAA. 52 p.
- Rabalais, N.N., R.E. Turner, B.K. Sen Gupta, D.F. Boesch, P. Chapman, M.C. Murrell. 2007. Hypoxia in the Northern Gulf of Mexico: Does the science support the plan to reduce, mitigate, and control hypoxia? *Estuaries and Coasts* 30:753-772.
- Rainer, S.F., R.C. Fitzhardinge. 1981. Benthic communities in an estuary with periodic deoxygenation. *Australian Journal of Marine and Freshwater Research* 32:227-243.
- Reish, D.J. 1955. The relation of polychaetous annelids to harbor pollution. *Public Health Report* 70:1168-1174.
- Reish, D.J. 2000. The seasonal settlement of polychaete larvae before and after pollution abatement in Los Angeles-Long Beach Harbors, California. *Bulletin of Marine Science* 67:672.
- Reizopoulou, S, M. Thessalou-Legaki, A. Nicolaidou. 1996. Assessment of disturbance in Mediterranean lagoons: An evaluation of methods. *Marine Biology* 125:189-197.
- Rhoads, J.M., D.J. Yozzo, M.M. Cianciola, R.J. Will. 2001. Norton Basin/Little Bay restoration project: historical and environmental background report. U.S. Army Corps of Engineers, New York District, CENAN-PL-ES, New York. p. 50.
- Riedel, G.F., J.G. Sanders, R.W. Osman. 1999. Biogeochemical control on the flux of trace elements from estuarine sediments: effects of seasonal and short-term hypoxia. *Marine Environmental Research* 47:349-372.
- Rikard, F.S., R.K. Wallace, D. Rouse, I. Saoud. 2000. The effect of low oxygen on oyster survival during reef restoration efforts in Bon Secour Bay, Alabama. *Journal of Shellfish Research* 19:640.
- Ritter, C., P.A. Montagna. 1999. Seasonal hypoxia and models of benthic response in a Texas bay. *Estuaries* 22:7-20.
- Rosenberg, R. 1980. Effects of oxygen deficiency on benthic macrofauna in fjords. *In: Fjord oceanography*, H.J. Freeland, D.M. Farmer and C.D. Levings (eds.), Plenum Publishing Corp., New York. pp. 499-514.
- Rosenberg, R. 1990. Negative oxygen trends in Swedish coastal bottom waters. *Marine Pollution Bulletin* 21:335-339.
- Rosenberg, R., R.J. Diaz. 1993. Sulphur bacteria (*Beggiatoa* spp.) mats indicate hypoxic conditions in the Inner Stockholm Archipelago. *Ambio* 22:32-36.
- Rosenberg, R., L.O. Loo. 1988. Marine eutrophication induced oxygen deficiency: effects on soft bottom fauna, western Sweden. *Ophelia* 29:213-225.
- Rosenberg, R., I. Olsson, E. Olundh. 1977. Energy flow model of an oxygen-deficient estuary on the Swedish west coast. *Marine Biology* 42:99-107.
- Rosenberg, R., L.O. Loo, P. Möller. 1992. Hypoxia, salinity and temperature as structuring factors for marine benthic communities in a eutrophic area. *Netherlands Journal of Sea Research* 30:121-129.

- Rosenberg, R., H.C. Nilsson, R.J. Diaz. 2001. Response of benthic fauna and changing sediment redox profiles over a hypoxic gradient. *Estuarine, Coastal and Shelf Science* 53:343-350.
- Rosenberg, R., J.S. Gray, A.B. Josefson, T.H. Pearson. 1987. Petersen's benthic stations revisited. II. Is the Oslofjord and eastern Skagerrak enriched? *Journal of Experimental Marine Biology and Ecology* 105:219-251.
- Rumohr, H. 1986. Historical evidence for eutrophication processes in the Kieler Bucht (western Baltic). *Meeresforschung* 31:115-123.
- Rybarczyk, H., B. Elkaim, J.G. Wilson, N. Loquet. 1996. Eutrophication in the Baie de Somme: Consequences and impacts on the benthic population. *Oceanologica Acta* 19:131-140.
- Sagasti, A., L.C. Schaffner, J.E. Duffy. 2001. Effects of periodic hypoxia on mortality, feeding and predation in an estuarine epifaunal community. *Journal of Experimental Marine Biology and Ecology* 258:257-283.
- Saiz-Salinas, J.I. G.Frances-Zubillaga. 1997. Tidal zonation on mud flats in a polluted estuary caused by oxygen-depleted water. *Journal of Experimental Marine Biology and Ecology* 209:157-170.
- Sale, J.W., W.W. Skinner. 1917. The vertical distribution of dissolved oxygen and the precipitation of salt water in certain tidal areas. *Franklin Institute Journal* 184:837-848.
- Sanger, D.M., M.D. Arendt, Y. Chen, E.L. Wenner, A.F. Holland, D. Edwards, J. Caffrey. 2002. A synthesis of water quality data: National Estuarine Research Reserve System-wide monitoring program (1995-2000). National Estuarine Research Reserve Technical Report Series 2002:3. South Carolina Department of Natural Resources, Marine Resources Division Contribution No. 500. 135 p.
- Santos, S.L., J.L. Simon. 1980. Marine soft-bottom community establishment following annual defaunation: larval or adult recruitment. *Marine Ecology Progress Series* 2:235-241.
- Saunders, J.E., K. M. A. Zahed, D. M. Paterson. 2007. The impact of organic pollution on the macrobenthic fauna of Dubai Creek (UAE). *Marine Pollution Bulletin* 54:1715-1723.
- Sanukida, S., H. Okamoto, M. Hitomi. 1984. Alteration of pollution indicator species of macrobenthos during stagnant period in eastern Hiuchi Sound. *Bulletin Japanese Society of Scientific Fisheries* 50:727.
- Savage, C., R. Elmgren, U. Larsson. 2002. Effects of sewage-derived nutrients on an estuarine macrobenthic community. *Marine Ecology-Progress Series* 243:67-82.
- Schimmel, S.C., S.J. Benyi, C.J. Strobel. 1999. An assessment of the ecological condition of Long Island Sound, 1990-1993. *Environmental Monitoring and Assessment* 56:27-49.
- Seliger, H.H., J.A. Boggs, W.H. Biggley. 1985. Catastrophic anoxia in the Chesapeake Bay in 1984. *Science* 228:70-73.
- Sen Gupta, B.K., R.E. Turner, N.N. Rabalais. 1996. Seasonal oxygen depletion in the continental shelf waters of Louisiana: Historical record of benthic foraminifers. *Geology* 24:227-230.
- Silke, J., F. O'Beirn, M. Cronin. 2005. *Karenia mikimotoi*: an exceptional dinoflagellate bloom in western Irish waters, summer 2005. *Marine Environment and Health Series*, No 21. Marine Institute, Marine Environment and Food Safety Services: Galway. p. 48.
- Sindermann, C., and R. Swanson. 1980. Chapter 1. Historical and Regional Perspective. *In*: Oxygen depletion and associated benthic mortalities in New York Bight, 1976. R.L. Swanson and C.J. Sindermann (eds.) Rockville. U.S. Department of Commerce National Oceanic and Atmospheric Administration. pp.1-16.

- Simunovic, A., C. Piccinetti, M. Zore-Armanda. 1999. Kill of benthic organisms as a response to an anoxic state in the northern Adriatic (a critical review). *Acta Adriatica* 40:37-64.
- Somerfield, P.J., V.G. Fonsêca-Genevois, A.C.L. Rodrigues, F.J.V. Castro, G.A.P. Santos. 2003. Factors affecting meiofaunal community structure in the Pina Basin, an urbanized embayment on the coast of Pernambuco, Brazil. *Journal of the Marine Biological Association of the United Kingdom* 83:1209-1213.
- Souchu, P., A. Gasc, Y. Collos, A. Vaquer, H. Tournier, B. Bibent, J-M. eslous-Paoli. 1998. Biogeochemical aspects of bottom anoxia in a Mediterranean lagoon (Thau, France). *Marine Ecology Progress Series* 164:135-146.
- Stachowitsch, M. 1984. Mass mortality in the Gulf of Trieste: The course of community destruction. *Marine Ecology* 5:243-264.
- Stachowitsch, M. 1991. Anoxia in the Northern Adriatic Sea: rapid death, slow recovery. *In: Modern and ancient continental shelf anoxia*. R.V. Tyson and T.H. Pearson (eds.). London: Geological Society Special Publication No. 58. pp. 119-129.
- Stanley, D.W., S.W. Nixon. 1992. Stratification and bottom-water hypoxia in the Pamlico River Estuary. *Estuaries* 15:270-281.
- Stora, G., A. Arnoux. 1983. Effects of large freshwater diversions on benthos of a Mediterranean lagoon. *Estuaries* 6:115-125.
- Summers, J.K., S.B. Weisberg, A.F. Holland, J. Kou, V.D. Engle, D.L. Breitberg, R.J. Diaz. 1997. Characterizing dissolved oxygen conditions in estuaries environments. *Environmental Monitoring and Assessment* 45:319-328.
- Suzuki, T. 2001. Oxygen-deficient waters along the Japanese coast and their effects upon the estuarine ecosystem. *Journal of Environmental Quality* 30:291-302.
- Suzuki, T., and Y. Matsukawa. 1987. Hydrography and budget of dissolved total nitrogen and dissolved oxygen in the stratified season in Mikawa Bay, Japan. *Journal of the Oceanographic Society of Japan* 43:37-48.
- Tagliapietra, D., M. Pavan, C. Wagner. 1998. Macrobenthic community changes related to eutrophication in Palude della Rosa (Venetian Lagoon, Italy). *Estuarine, Coastal and Shelf Science* 47:217-226.
- Taylor, D. I. 2000. Harbor sampling update. MWRA Sewerage Division. Online at: <http://www.mwra.state.ma.us/harbor/html/wk090100.htm>
- Taylor, F.J., N.J. Taylor, J.R. Walsby. 1985. A bloom of the planktonic diatom, *Cerataulina pelagica*, off the coast of northeastern New Zealand in 1983, and its contribution to an associated mortality of fish and benthic fauna. *Int Revue ges Hydrobiologia* 70:773-795.
- Tenore, K.R. 1972. Macrobenthos of the Pamlico River estuary, North Carolina. *Ecological Monographs* 42:51-69.
- Theodorou, A.J. 1996. Long-term environmental effects of raw sewage sea disposal in Elefsis Bay (Saronikos Gulf, Greece). *Partnership in Coastal Zone Management*, Samara Publishing Ltd., Cardigan, UK. pp. 547-556.
- Thiel, R., A. Sepulveda, R. Kafemann, W. Nellen. 1995. Environmental factors structuring the fish community of the Elbe estuary. *Journal of Fisheries Biology* 46:47-69.
- Thomassin, B.A., P. Gourbesville, B. Gout, A. Arnoux. 1998. Impact of an industrial and urban sewage off a coral fringing reef at Mauritius (Indian ocean): modeling of plumes, distribution of trace metals in sediments and effects of the eutrophication on coral reef communities. *Oceans 98 - Engineering for sustainable use of the oceans: conference proceedings, Nice, IEEE/OES, Piscataway, New Jersey*. 1:301-305

- Thouvenin, B., P. Le Hir, L.A. Romana. 1994. Dissolved oxygen model in the Loire Estuary. Changes in fluxes in estuaries: implications from science to management. Olsen and olsen, Fredensborg, Denmark. pp. 169-178.
- Tomasko, D.A., C. Anastasiou, C. Kovach. 2006. Dissolved oxygen dynamics in Charlotte Harbor and its contributing watershed, in response to hurricanes Charley, Frances, and Jeanne - impacts and recovery. *Estuaries and Coasts* 29:932-938.
- Toner, P., J. Bowman, K. Clabby, J. Lucey, M. McGarrigle, C. Concannon, C. Clenagh, J. Delaney, S. O'Boyle, M. MacCa'rthaigh, M. Craig, R. Quinn. 2005. Water quality in Ireland 2001-2003. Environmental Protection Agency, County Wexford, Ireland. 232 p.
- Tsutsumi, H. 1987. Population dynamics of *Capitella capitata* (Polychaeta, Capitellidae) in an organically polluted cove. *Marine Ecology Progress Series* 36:139-149.
- Tulkki, P. 1965. Disappearance of the benthic fauna from the basin of the Bornholm (southern Baltic) due to oxygen deficiency. *Cahiers de Biologie Marine* 6:455-463.
- Tunncliffe, V. 1981. High species diversity and abundance of the epibenthic community in an oxygen-deficient basin. *Nature* 294:354-356.
- Turner, R.E., N.N. Rabalais. B. Fry, N. Atilla, C.S. Milan, J.M. Lee, C. Normandeau, T.A. Oswald, E.M. Swenson, D.A. Tomasko. 1994. Paleo-indicators and water quality change in the Charlotte Harbor estuary (Florida). *Limnology and Oceanography* 51:518-523.
- Turner, R.E., N.N. Rabalais. 1994. Coastal eutrophication near the Mississippi river delta. *Nature* 368:619-621.
- Tyler, R.M. 2004. Distribution and avoidance patterns of juvenile summer flounder (*Paralichthys dentatus*) and weakfish (*Cynoscion regalis*) in relation to hypoxia: field studies in a temperate coastal lagoon tributary and laboratory choice-trial experiments. PhD thesis, University of Delaware, Newark, DE.
- Ueda, N., H. Tsutsumi, M. Yamada, K. Hanamoto, S. Montani. 2000. Impacts of oxygen-deficient water on the macrobenthic fauna of Dokai Bay and on adjacent intertidal flats, in Kitakyushu, Japan. *Marine Pollution Bulletin* 40:906-913.
- Ukwe, C.N., C.A. Ibe, P.C. Nwilo, P.A. Huidobro. 2006. Contributing to the WSSD targets on oceans and coasts in west and central Africa: the Guinea current large marine ecosystem project. *International Journal of Oceans and Oceanography* 1:21-44.
- Uncles, R.J., R.J.M. Howland, A.E. Easton, M.L. Griffiths, C. Harris, R.S. King, A.W. Morris, D.H. Plummer, E.M.S. Woodward. 1998. Seasonal variability of dissolved nutrients in the Humber-Ouse Estuary, UK. *Marine Pollution Bulletin* 37:234-246.
- Valdes-Lozano, D.S., M. Chumacero, E. Real. 2006. Sediment oxygen consumption in a developed coastal lagoon of the Mexican Caribbean. *Indian Journal of Marine Sciences* 35:227-234.
- Valentin, J., D. Tenenbaum, A. Bonecker, S. Bonecker, C. Nogueira, R. Paranhos, M.C. Villac. 1999. Hydrobiological characteristics of the Guanabara bay (Brazil). *Journal de Recherche Océanographique* 24:33-41.
- Velasquez, I.B., G.S. Jacinto, F.S. Valera. 2002. The speciation of dissolved copper, cadmium and zinc in Manila Bay, Philippines. *Marine Pollution Bulletin* 45:210-217.
- Verlaan, P.A.J., M. Donze, P. Kuik. 1998. Marine vs fluvial suspended matter in the Scheldt Estuary. *Estuarine, Coastal and Shelf Science* 46:873-883.

- Wang, B. 2006. Cultural eutrophication in the Changjiang (Yangtze River) plume: history and perspective. *Estuarine, Coastal and Shelf Science* 69:471-477.
- Wang, T. 2005. Hypoxia in shallow coastal waters: a case study in Onancock Creek, Virginia. Masters of Science Thesis, Virginia Institute of Marine Science, College of William and Mary, Gloucester Pt. 129 p.
- Ward, G.H., Jr. 2000. Effects of shrimp farming on the hydrography and water quality of El Pedregal and San Bernardo Estuaries, Gulf of Fonseca, Honduras. CRSP Research Report 00-150:1-32.
- Weigelt, M. 1990. Oxygen conditions in the deep water of Keil Bay and the impact of inflowing salt-rich water from the Kattegatt. *Meeresforschung* 33:1-22.
- Weigelt, M. 1991. Short- and long-term changes in the benthic community of the deeper parts of Kiel Bay (Western Baltic) due to oxygen depletion and eutrophication. *Meeresforschung* 33:197-224.
- Weisberg, S.B., H.T. Wilson, P. Himchak, T. Baum, R. Allen. 1996. Temporal trends in abundance of fish in the tidal Delaware River. *Estuaries* 19:723-729.
- Welsh, B., R. Welsh, M. DiGiacomo-Cohen. 1994. Quantifying hypoxia and anoxia in Long Island Sound. *In: Changes in fluxes in estuaries: Implications from science to management.* K. Dyer, and R. Orth (eds.) Fredensborg: Olsen and Olsen. pp. 131-137.
- Westernhagen, H.V., V. Dethlefsen. 1983. North Sea oxygen deficiency 1982 and its effects on the bottom fauna. *Ambio* 12:264-266.
- Wilkin, R., H.L. Barnes. 1997. Pyrite formation in an anoxic estuarine basin. *American Journal of Science* 297:620-650.
- Wu, R.S.S. 1982. Periodic defaunation and recovery in a subtropical epibenthic community, in relation to organic pollution. *Journal of Experimental Marine Biology and Ecology* 64:253-269.
- Yamamuro, M., J. Hiratsuka, Y. Ishitobi. 2000. Seasonal change in a filter-feeding bivalve *Musculista senhousia* population of a eutrophic estuarine lagoon. *Journal of Marine Systems* 26:117-126.
- Yamamuro, M., N. Oka, J. Hiratsuka. 1998. Predation by diving ducks on the biofouling mussel *Musculista senhousia* in a eutrophic estuarine lagoon. *Marine Ecology Progress Series* 174:101-106.
- Yokoyama, H. 1995. Occurrence of *Paraprionospio* sp. (Form A) larvae (Polychaeta: Spionidae) in hypoxic water of an enclosed bay. *Estuarine, Coastal and Shelf Science* 40:9-19.
- Yokoyama, H. 2002. Impact of fish and pearl farming on the benthic environments in Gokasho Bay: evaluation from seasonal fluctuations of the macrobenthos. *Fisheries Science* 68:258-268.
- Zaitsev, Y.P. 1993. Impacts of eutrophication on the Black Sea fauna. *In: Fisheries and environment studies in the Black Sea system,* Food and Agricultural Organization of the United Nations, Rome, Italy. pp. 64-86.
- Zimmerman, A.R., E.A. Canuel. 2000. A geochemical record of eutrophication and anoxia in Chesapeake Bay sediments: anthropogenic influence on organic matter composition. *Marine Chemistry* 69:117-137.